



US006723549B2

(12) **United States Patent**
Miettinen-Oinonen et al.

(10) **Patent No.:** **US 6,723,549 B2**
 (45) **Date of Patent:** ***Apr. 20, 2004**

(54) **CELLULASES, THE GENES ENCODING THEM AND USES THEREOF**

5,525,507 A 6/1996 Clarkson et al. 435/263
 5,650,322 A 7/1997 Clarkson et al. 435/263
 6,001,639 A 12/1999 Schulein et al. 435/263

(75) **Inventors:** Arja Miettinen-Oinonen, Masala (FI);
 John Londesborough, Helsinki (FI);
 Jari Vehmaanperä, Klaukkala (FI);
 Heli Haakana, Rajamäki (FI); Arja
 Mäntylä, Helsinki (FI); Raija Lanto,
 Klaukkala (FI); Minna Elovainio,
 Helsinki (FI); Vesa Joutsjoki, Helsinki
 (FI); Marja Paloheimo, Vantaa (FI);
 Pirkko Suominen, Helsinki (FI)

(73) **Assignee:** AB Enzymes Oy (FI)

(*) **Notice:** This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** 08/841,636

(22) **Filed:** Apr. 30, 1997

(65) **Prior Publication Data**

US 2002/0168751 A1 Nov. 14, 2002

Related U.S. Application Data

(63) Continuation of application No. PCT/FI96/00550, filed on Oct. 17, 1996, and a continuation-in-part of application No. 08/732,181, filed on Oct. 16, 1996.

(60) Provisional application No. 60/005,335, filed on Oct. 17, 1995, provisional application No. 60/007,926, filed on Dec. 4, 1995, and provisional application No. 60/020,840, filed on Jun. 28, 1996.

(51) **Int. Cl.⁷** C12N 9/42; C11D 3/386

(52) **U.S. Cl.** 435/204; 435/263; 510/392

(58) **Field of Search** 475/209, 263; 510/392

FOREIGN PATENT DOCUMENTS

DE	1 767 737	9/1971
EP	0 220 016	4/1987
EP	0 238 216	9/1987
EP	0 307 564	3/1989
EP	0 383 828	8/1990
EP	0 406 314	1/1991
EP	0 510 091	10/1992
EP	0 521 999	1/1993
EP	0 531 372	3/1993
EP	0 576 526	1/1994
GB	2 075 028	11/1981
WO	WO 89/09259	10/1989
WO	WO 91/14822	10/1991
WO	WO 91/17243	11/1991
WO	WO 91/17244	11/1991
WO	WO 92/06165	4/1992
WO	WO 92/06209	4/1992
WO	WO 92/06210	4/1992
WO	WO 92/06221	4/1992
WO	WO 92/17572	10/1992
WO	WO 92/17574	10/1992
WO	WO 92/18599	10/1992
WO	WO 93/11249	6/1993
WO	WO 93/22428	11/1993
WO	WO 93/24621	12/1993
WO	WO 93/24622	12/1993
WO	WO 94/07983	4/1994
WO	WO 94/07998	4/1994
WO	WO 94/14953	7/1994
WO	WO 94/23113	10/1994
WO	WO 95/02675	1/1995
WO	WO 95/16782	6/1995
WO	WO 95/24471	9/1995
WO	WO 95/25840	9/1995
WO	WO 96/17994	6/1996
WO	WO 96/29397	9/1996
WO	WO 96/34945	11/1996

OTHER PUBLICATIONS

Sequence search, 1999.*

(List continued on next page.)

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,438,864 A	4/1969	Tendler	195/66
3,664,961 A	5/1972	Norris	252/99
4,106,989 A	8/1978	Komura et al.	195/62
4,832,864 A	5/1989	Olson	252/174.12
4,894,338 A	1/1990	Knowles et al.	435/172.3
4,912,056 A	3/1990	Olson	435/263
5,006,126 A	4/1991	Olson et al.	8/401
5,120,463 A	6/1992	Bjork et al.	252/174.12
5,122,159 A	6/1992	Olson et al.	8/401
5,213,581 A	5/1993	Olson et al.	8/401
5,232,851 A	8/1993	Cox et al.	435/263
5,246,853 A	9/1993	Clarkson et al.	435/263
5,290,474 A	3/1994	Clarkson et al.	252/174.12
5,298,405 A	3/1994	Nevalainen et al.	435/209
5,443,750 A	8/1995	Convents et al.	252/174.12
5,520,838 A	5/1996	Baeck et al.	252/174.12

Primary Examiner—Charles L. Patterson, Jr.

(74) *Attorney, Agent, or Firm*—Sterne, Kessler, Goldstein & Fox P.L.L.C.

(57) **ABSTRACT**

Genes encoding novel cellulases, and a gene encoding a protein that facilitates the action of such novel cellulases, the novel cellulases and a protein that facilitates the action of such cellulases, and enzyme preparations containing such proteins are described. The native hosts and the culture medium of said hosts containing said novel cellulases are also disclosed. These proteins are especially useful in the textile and detergent industry and in pulp and paper industry.

16 Claims, 37 Drawing Sheets

OTHER PUBLICATIONS

- Tomme, P. et al. (1988) *Eur. J. Biochem.* 170:575-581.
- Offord, D. A., et al. (1991) *Appl. Biochem. Biotechnol.* 28/29, 377-385.
- Aho, S. et al., "Monoclonal antibodies against core and cellulose-binding domains of *Trichoderma reesei* cellobiohydrolases I and II and endoglucanase I," *Eur. J. Biochem.* 200:643-649 (1991).
- Bailey, M.J. and Nevalainen, K.M.H., "Induction, isolation and testing of stable *Trichoderma reesei* mutants with improved production of solubilizing cellulase," *Enz. Micr. Tech.* 3(2):153-157 (1981).
- Bhat, K.M. et al., "The endo-(1→4)-β-D-glucanase system of *Penicillium pinophilum* cellulase: isolation, purification, and characterization of five major endoglucanase components," *Carbohydrate Research* 190:279-297 (1989).
- Bhikhabhai, R. et al., "Isolation of Cellulolytic Enzymes from *Trichoderma reesei* QM 9414," *J. Applied Biochem.* 6:336-345 (1984).
- Bühler, R., "Double-Antibody Sandwich Enzyme-Linked Immunosorbent Assay for Quantitation of Endoglucanase I of *Trichoderma reesei*," *Appl. Environ. Microbiol.* 57(11):3317-3321 (1991).
- Davies, G. J. et al., "Structure and function of endoglucanase V," *Nature* 365(6444):362-364 (1993).
- Harkki, A. et al., "Genetic engineering of *Trichoderma* to produce strains with novel cellulase profiles," *Enzyme Microb. Technol.* 13(3):227-233 (1991).
- Henrissat, B. and Bairoch, A., "New families in the classification of glycosyl hydrolases based on amino acid sequence similarities," *Biochem.J.* 293:781-788 (1993).
- Karhunen, T. et al., "High frequency one-step gene replacement in *Trichoderma reesei*. I. Endoglucanase I overproduction," *Mol. Gen. Genet.* 241:515-522 (1993).
- Klahorst, S. et al., "Optimizing the Use of Cellulase Enzymes," *Textile Chemist and Colorist* 26(2):13-18 (Feb. 1994).
- Klahorst, S. et al., "Optimizing the Use of Cellulases for Denim Finishing," *AATCC Int. Conf. & Exhibit*, Atlanta GA, Oct. 4-7, 1992, p. 243.
- Kochavi, D. et al., "Optimizing Processing Conditions In Enzymatic Stonewashing," *American Dyestuff Reporter* 79(9):24, 26, 28 (1990).
- Lange, N.K., "Application of cellulases in the textile industry," *Proc. Second TRICEL Symp. On Trichoderma reesei Cellulases and Other Hydrolases*, Espoo Finland, 1993, P. Suominen et al. ed., Foundation for Biotechnical and Industrial Fermentation Research 8 (1993):263-272.
- Lantto, R. et al., "Backstaining In Denim Wash With Different Cellulases," *Amer. Dyestuff Reporter*, pp. 64-65, and 72 (Aug. 1996).
- Mach, R.L. et al., "Transformation of *Trichoderma reesei* based on hygromycin B resistance using homologous expression signals," *Curr. Genet.* 25(6):567-570 (Jun. 1994).
- Maloney, A.P. et al., "Cellulose Hydrolysis by the Cellulases Produced by *Talaromyces emersonii* when Grown on Different Inducing Substrates," *Biotechnol. Bioeng.* 25(4):1169-1173 (1983).
- Mandels, M. and Reese, E. T., "Induction of cellulase in *Trichoderma viride* as influenced by carbon sources and metals," *J. Bacteriol.* 73(2):269-278 (1957).
- Mattern, I. E. et al., "Transformation of *Aspergillus Oryzae*," in Abstracts of the 19th Lanteren Lectures on Molecular Genetics of Yeasts and Filamentous Fungi and its Impact on Biotechnology, Lanteren, the Netherlands, 1987, p. 34.
- Miettinen-Oinonen, A. et al., "Effects of Cellulases on Cotton Fiber and Fabric," *Proceedings of the 77th World Conference of the Textile Institute* 1(2):197-207 (May 1996).
- Nevalainen, H., "Genetic improvement of enzyme production in industrially important fungal strains," VTT, Technical Research Centre of Finland, publication 26, Espoo, Finland (1985).
- Nilsson, T.E., "Enzymes for the Textile Industry--Traditional and New Applications," presented at Aachen Textile Conference, Nov. 22-23, 1994, *DWI Reports* 114:85-88 (Feb. 1995).
- Penttilä, M. et al., "Homology between cellulase genes of *Trichoderma reesei*: complete nucleotide sequence of the endoglucanase I gene," *Gene* 45:253-263 (1986).
- Penttilä, M. et al., "A versatile transformation system for the cellulolytic filamentous fungus *Trichoderma reesei*," *Gene* 61:155-164 (1987).
- Saloheimo, A. et al., "A novel, small endoglucanase gene, eg15, from *Trichoderma reesei* isolated by expression in yeast," *Molec. Microbiol.* 13(2):219-228 (Jul. 1994).
- Saloheimo, M. et al., "EGIII, a new endoglucanase from *Trichoderma reesei*: the characterization of both gene and enzyme," *Gene* 63:11-21 (1988).
- Schindler, M. et al., "Characterization of the pyruvate kinase-encoding gene (pkil) of *Trichoderma reesei*," *Gene* 130:271-275 (1993).
- Schüle, M., "Cellulases of *Trichoderma reesei*," *Methods Enzymol.* 160:234-242 (1988).
- Schüle, M. et al., "Humicola insolens Alkaline Cellulases," *Proceedings of the second TRICEL symposium on Trichoderma reesei Cellulases and Other Hydrolases*, Espoo 1993, P. Suominen and T. Reinikainen eds., Foundation for Biotechnical and Industrial Fermentation Research 8:109-116 (1993).
- Shoemaker, S. et al., "Molecular cloning of exo-cellobiohydrolase I derived from *Trichoderma reesei* strain L27," *Bio/Technology* 1:691-696 (1983).
- Suominen, P. et al., "High frequency one-step gene replacement in *Trichoderma reesei*. II. Effects of deletions of individual cellulase genes," *Mol. Gen. Genet.* 241:523-530 (1993).
- Teeri, T. T. et al., "The cellulolytic enzyme system of *Trichoderma reesei*: Molecular cloning, characterization and expression of the cellobiohydrolase genes," *Technical Research Center of Finland, Espoo 1987, Publications* 38.
- Teeri, T. et al., "Homologous domains in *Trichoderma reesei* cellulolytic enzymes: gene sequence and expression of cellobiohydrolase II," *Gene* 51:43-52 (1987).
- Teeri, T. et al., "The molecular cloning of the major cellulase gene from *Trichoderma reesei*," *Bio/Technology* 1:696-699 (1983).
- Tyndall, R. Michael, "Application of Cellulase Enzymes to Cotton Fabrics and Garments," *AATCC Book of Papers: 1991 International Conference & Exhibition*, American Association of Textile Chemists and Colorists, Research Triangle Park, N.C., pp. 269-273 (1991).

- Tyndall, R. Michael, "Improving the Softness and Surface Appearance of Cotton Fabrics and Garments by Treatment with Cellulase Enzymes," *Textile Chemist and Colorist* 24(6):23-26 (1992).
- van Tilbeurgh, H. et al., "Fluorogenic and Chromogenic Glycosides as Substrates and Ligands of Carbohydrases," *Methods Enzymol.* 160:45-59 (1988).
- Videbaek, T. et al., "The jeans effect comes into being," *ITB Dyeing/Printing/Finishing* (Jan. 1994) pp. 25-29.
- Wood, T.M. et al., "Aerobic and Anaerobic Fungal Cellulases, with Special Reference to Their Mode of Attack on Crystalline Cellulose," *FEMS Symposium No. 43: Biochemistry and Genetics of Cellulose Degradation*, pp. 31-52 (1988).
- "Celluzyme™" Product Sheet, NOVO Detergent Enzymes Division, Novo Nordisk A/S (date unknown).
- "Denimax™ T" Product Sheet, Enzyme Process Division, Novo Nordisk (date unknown).
- "Denimax™ L" Product Sheet, Enzyme Process Division, Novo Nordisk (date unknown).
- "Denimax™ PB" Product Sheet, Enzyme Process Division, Novo Nordisk (date unknown).
- "Denimax™ BT" Product Sheet, Enzyme Process Division, Novo Nordisk (date unknown).
- Measurement of Cellulase Activities, Commission on Biotechnology: International Union of Pure and Applied Chemistry, Biochemical Engineering Research Centre, Indian Institute of Technology, Delhi, New Delhi-110016, pp. 1-13 (1984).
- International Search Report for International Application No. PCT/FO96/00550, mailed Feb. 14, 1997.
- El-Gindy, A.A., "Production of Cellulase(s) by *Myriococcum albomyces*," *Zentralbl. Mikrobiol.* 146:193-96 (1991).
- Latif F., "Saccharification of *Leptochloa fusca* (Kallar Grass Straw) Using Thermostable Cellulases," *Bioresource Technol.* 50:107-111 (1994).
- Biotechnology Abstracts AN 91-08942 Abstract for El-Gindy, A.A., *Zentralbl. Mikrobiol.* 146:193-96 (1991) (Document AT16).
- Biotechnology Abstracts AN 95-04270 Abstract for Latif, F. et al., *Bioresource Technol.* 50:107-111 (1994) (Document AT16).

* cited by examiner


ALK04179: pH-DEPENDENCY OF
THE ENDOGLUCANASE ACTIVITY

FIG.1A


ALK04179: TEMPERATURE DEPENDENCY OF
THE ENDOGLUCANASE ACTIVITY

FIG.1B


ALKO4124: pH-DEPENDENCY OF
THE ENDOGLUCANASE ACTIVITY

FIG.2A


ALKO4124: TEMPERATURE DEPENDENCY OF
THE ENDOGLUCANASE ACTIVITY

FIG.2B




FIG.3A




FIG.3B




FIG.4A




FIG.4B




FIG.5A




FIG.5B




FIG.6A




FIG.6B




FIG.7A

BLUENESS WITH ECOSTONE L




FIG.7B




FIG. 8




FIG.9A




FIG.9B




FIG.10




FIG.11A




FIG.11B

TEMPERATURE DEPENDENCE OF THE
ENDOGLUCANASE ACTIVITY OF 50K-CELLULOSE
AT pH 7.0




FIG.12

THE pH DEPENDENCE OF THE ENDOGLUCANASE
ACTIVITY OF 50K-CELLULOSE AT 50°C AND 70°C




FIG.13




FIG.14

20K-CELLULASE: TEMPERATURE DEPENDENCE OF
ENDOGLUCANASE ACTIVITY AT pH 7 (10 MIN REACTION
TIMES)




FIG.15

20K-CELLULASE: pH-DEPENDENCE OF ENDOGLUCANASE
ACTIVITY (10 MIN REACTION TIMES)




FIG.16A

20K-CELLULASE: pH-DEPENDENCE OF ENDOGLUCANASE
ACTIVITY (60 MIN REACTION TIMES)




FIG.16B

429
ANGQSTRYWDCCCKPSCGWRGKGPVNQPVYS

430
YGGISSR

431
CGWR

432
PSCGWR

433
YWDCCCK

439
QECDSFPEPLKPGCQWR

fr 9
RHDDGGFA

fr 14
YWDCCCKP

fr 16
GKGPVNQPVYSCDANFQR

fr 17
VQCPEELVAR

fr 28
DWFQNADNPSFTFER

fr 30
TMVQSTSTGGDLGSNHFDLNIPGGGVGLF

FIG.17




FIG.18


-30 -10 10 30 50
TGGCCCTAACCGAGAACCAAGACTCCAAGAATGGCTCTACTCCCGTTCTCGCGGCCCTCTCTGGCCGGAGCATTTGCCCTCGGGCCCTCGCCGCCAA
M R S T P V L R A L L A A L P L G A L A A N
70 90 110 130 150
CGGTCACTCCACGAGGtaactgatcacccgcctcattacgcgtgcgcgaccgcggttcagggtcactgctcacccgcatccagATACTGGGACTGCT
G Q S T R Y W D C C
170 190 210 230 250
GCAAGCCGTCTGTGGGCTGGCGCGGAAGGGCCCGGTGAACCAAGCCCGTCTACTCGTGGAGCGCAACTTCCAGCGCATCCACGACTTCGATGCCGCTCTC
K P S C G W R G K G P V N Q P V Y S C D A N F Q R I H D F D A V S
270 290 310 330 350
GGGTGCGAGGGGCGCCCGCCTTCTGTGTGGCGGACCAACAGCCCTGGGCCATTATGACAACCTCTGTACGGCTTCGGGGGCGACTGCACTCAGCGGC
G C E G G P A F S C A D H S P W A I N D N L S Y G F A A T A L S G
370 390 410 430 450
CAGACCGAGGAGTCGTGGTGTCTGTGCTGTCTACGCGtgtagtgcttggcccaacgtcggtgattccggaggttcagaccactgaccagcagcccgctc
Q T E E S W C C A C Y A

FIG.19A

890

CGTGTGACGGCAGCTACGTGGAACGACCTGGAGCTC

Fig. 19B



-230 -210 -190 -170 -150
GAATTGGGGGTTGCCAGGGAGTCGTACAGGGGTGGGTGGAGGGGGATGGGGATGGAGAAGCAATATATGGAGCTTTGTGCTC
-130 -110 -90 -70 -50
GGGGCTCCCTCTGCCACGTTCCCTTGCCCTTGCCTGGGTGTTGTGGTCTTCCCTTACCATCCGACAAACCAACCTGCTGCGGGTGAACCTCGCA
-30 -10 10 30 50
GAGCGCCTTCGGACGACGACAGACAGCGCACCATGACTCGCAACATCGCCCTGCTCGGGCGCGCGTGGGGCTCCTGGGCCCTGCGCCACGGCCAGAAGC
M T R N I A L L G A A S A L L G L A H G Q K P
70 90 110 130 150
CGGGCGAGACGCCCGAGGTGCACCCGCAAGCTGACGACGTTCCGGTGACCAAGCGGACGGGTGCCAGCCGGGACCAACTACATTGTGCTGGACTCGCT
G E T P E V H P Q L T T F R C T K A D G C Q P R T N Y I V L D S L
170 190 210 230 250
GTGCAACCCGGTGCACCAAGTGGACAACGACTACAACTGCGGCGGACTGGGGGCAGAAAGCCCAACGGACGGCGTGCCTGGGACGCTCGAGTCGTGCGCGCGC
S H P V H Q V D N D Y N C G D W G Q K P N A T A C P D V E S C A R
270 290 310 330 350
AACTGCATCATGGAGGGGTGCCCGACTACAGCCAGCAGCGCGTCCAGCAGCGGACACGTCGCTGGGCCCTGCAGCAGCTCGTCGACGGCCCGCCTCGTCA
N C I M E G V P D Y S Q H G V T T S D T S L R L Q Q L V D G R L V T

FIG.21A

370 390 410 430 450
CGCGCGCGTCTACCTGCTGACGAGACCGAGCACGCTACGAGATGATGCACCTGACCGGCCAGGAGTTACCTTTGAGGTGCGACGCCACCAAGTGCC
P R V Y L L D E T E H R Y E M H L T G Q E F T F E V D A T K L P

470 490 510 530 550
CTGGGCATGAACAGCGCCCTCTACCTGTCTCGAGATGGACCCGACCGCGGCCGAGCGAGCTCAACCCCGCGGTGCCTACTACGGCACCGGGCTACTGC
C G M N S A L Y L S E M D P T G A R S E L N P G G A Y Y G T G Y C

570 590 610 630 650
GAGCCCCAGTGTCTCGTGACGCCATTTCATCAACGGCATTgtgagtgttccctttggccccccctgaaatagatgtacctggtgtaacccccggg
D A Q C F V T P F I N G I

670 690 710 730 750
tgtcgcaccaaaacagGGCAACATCGAGGGCAAGGGCTCGTGTGCAACGAGATGGACATCTGGGAGGCCAACTCGCGGGCGACGACGTGGCGCGCAC
G N I E G G K G S C C N E M D I W E A N S R A T H V A P H

770 790 810 830 850
ACGTGCAACAGACGGGTCTGTACATGTGCGAGGGCGCGAGTACGACGGGTGTGCGACAAGGACGGGTGCGGGTGGAAACCCGTACCGGGTCA
T C N Q T G L Y M C E G A E C E Y D G V C D K D G C G W N P Y R V N

870 890 910 930 950
ACATCACCAGCTACTACGGCAACTCGGACGGGTTCGGCGTCGACACGGCGGGCCCTTCACCGTGGTGACGCGAGTCCCGGCCGACCGCGAGGGCGGCT
I T D Y Y G N S D A F R V D T R R P F T V V T Q F P A D A E G R L

FIG. 21B

970 990 1010 1030 1050
CGAGAGCATCCACCGGCTGTACGTGCAGGACGGCAAGGTGATCGAGTCTACGTCTGTCGACGGCCGGGCTGCCCGGACCGACTCGCTCAACGACGAG
E S I H R L Y V Q D G K V I E S Y V V D A P G L P R T D S L N D E

1070 1090 1110 1130 1150
TTCTGCGCGCCACGGCGCGCGCTACCTCGACCTCGCGGCACCGGGCATGGGCGACGCCCATGACGCGGGCATGGTGTGGCCATGAGCATCT
F C A A T G A A R Y L D L G G T A G M G D A M T R G M V L A M S I W

1170 1190 1210 1230 1250
GGTGGGACGAGTCCGGCTTCATGAACCTGGCTCGACAGCGGCGAGCGCGGCCCTGCCTGCCCGACGAGGGGACCCCAAGAACATTGTCAAGGTCGAGCC
W D E S G F M N W L D S G E A G P C L P D E G D P K N I V K V E P

1270 1290 1310 1330 1350
CAGCCCCGAGGTCACTACAGCAACCTGCGCTGGGGCGAGATCGGGTCGACCTTTGAGGCCGAGTCCGACGACGACGCGGCGACCGACTGCTAGATA
S P E V T Y S N L R W G E I G S T F E A E S D D G D G D C *

1370 1390 1410 1430 1450
ACTAACTAGTGGCGGAAGGGCGGGGATGCGTAACCTTACATACAGCCGGAGTTGTTTGAGTGTAGAGTATTGAGCTTTTCGATGTTAGTTGAGTG

1470 1490 1510 1530 1550
GAA TGGA AAAATTGCGTCTTTGCCCGGTGGTTGCGATAAACAATAGTCGGCTGGTGCATTTGTGACACTTCAATTGCGCTGTTGGCTTGGTGACAGACA

1570 1590 1610 1630 1650
CGGCGCGTCGATGACCCGACACCCAGAAATAATTCGCATGGTTGATTANTGTATTGTGCTTTAAATCGGAGGCTGATGCTCATCTCTCGAAATTC

FIG. 21C




FIG.22

-150 -130 -110 -90 -70
CCCGGTCTGGAGACGGGGAGCGGCCAGCGACGAGGATAAGAAGGCGACGACCGCGCCTCCGAGCCAGGCCCCAGGACAGGAGAACTCGGCCACGGCG
-50 -30 -10 10 30
AAGCAGCACGCCCGGATCGACAGTGTCCGCTCTGCCACAGCACTCTGCAACCATGATGTAAGCAGTACCTCCAGTACCTCGCGGGCGCGCTGCCGCT
M M K Q Y L Q Y L G A C L P L
50 70 90 110 130
CGTCGGCCTCGCCGCGGCCAGCGCGTGGTAACGAGAGCGCCGAGAACCAACCCCGCTCACCTGGCAGAGGTGACGGGCCCGGGCAACTGCCAGACC
V G L A A G Q R A G N E T P E N H P P L T W Q R C T A P G N C Q T
150 170 190 210 230
GTGAACGCCGAGGTGCTCATTGACGCCAACTGGCGCTGGCTGCACGACGACACAACATGCAGAACTGCTACGAGGGCAACCAAGTGGACCAACGCCCTGCAGCA
V N A E V V I D A N W R W L H D D N M Q N C Y D G N Q W T N A C S T
250 270 290 310 330
CCGCCACCGACTGCGCTGAGAAGTGCATGATCGAGGGTGCCGGCGACTACCTGGGCACCTACGGCGCCTCGACCAGCGGGCGCCCTGACCGCTCAAGTT
A T D C A E K C M I E G A G D Y L G T Y G A S T S G D A L T L K F
350 370 390 410 430
CGTCACCAAGCAGCAGTACGGCACCACCAACGTGCGGCTCGGGCTTCTACCTCATGAACGGGCCCGGACAAGTACCAGATGTTCAACCTCATGGGGCAACGAGCTT
V T K H E Y G T N V G S R F Y L M N G P D K Y Q M F N L M G N E L

FIG.23A

450 470 490 510 530
GCCTTTGACGTGACCTCTCGACCGTCGAGTGGGCATCAACAGCGCCCTGTACTTCGTGCGCATGGAGGAGGACGGCGGCATGGCCAGCTACCCGAGCA
A F D V D L S T V E C G I N S A L Y F V A M E D G G M A S Y P S N
550 570 590 610 630
ACCAGCGCGCGCCCGGTACGGCAGTGGGtgagttgagctccgcttctgagtcgcaacgaggcactttctggcgccggcctaactctctcgattc
Q A G A R Y G T G
650 670 690 710 730
ctccgacagTACTGGATGCCCAATCGCGTCGTGATCTCAAGTTCGTGGCGGCAAGGCCAACATTGAGGGCTGGAAGTGCTCCACCGAGCGACCCCAACG
Y C D A Q C A R D L K F V G G K A N I E G W K S S T S D P N A
750 770 790 810 830
CTGGCTGGGCGCGTACGGCAGCTGCTGGCGTGAGATCGACGTCGTgagtcgagacccgtccaccagggttcggatgcgggttggaatttcgcggct
G V G P Y G S C C A E I D V W
850 870 890 910 930
aacggagcaccctccagGGAGTCGAATGCCTATGCCTTCGCTTTCAGCGCGCAGCGGTGCAGGACCAACGAGTACCAGTCTGCGAGAGACCACCAACTGCG
E S N A Y A F A F T P H A C T T N E Y H V C E T T N C G
950 970 990 1010 1030
GTGGCACCTACTGGAGGACCGCTTCGCGGCAAGTGGACGCCAAGGGCTGGACTACAAACCCCTACCGCATGGGCAACCCGACTTCTACGGCAAGGG
G T Y S E D R F A G K C D A N G C D Y N P Y R M G N P D F Y G K G
1050 1070 1090 1110 1130
CAAGACGCTCGACACCGCGCAAGTTCACgtgcgtgaccccttgaggcgaaccttctctgcctgacctggacacactgaaactgacacgtcggttttcg
K T L D T S R K F T

FIG. 23B

1150 1170 1190 1210 1230
gctgcagCGTcGtCtCCCGcTTCgAGGAGaACAAGcTCTCCAGtACTTcATcCAGGAGCGCCGCAAGATCGAGATCCCGCCGCGACGTGGGAGGGCAT
V V S R F E E N K L S Q Y F I Q D G R K I E I P P T W E G M

1250 1270 1290 1310 1330
GCCCAACAGcAGcGAGATcACCCcCGAGcTcTcGCTCCACCATgTTCGATgTGTCAACGACCGCAACCGcTTCGAGGAGGTTCGGCGGcTTCGAGcAGCTG
P N S S E I T P E L C S T M F D V F N D R N R F E E V G G F E Q L

1350 1370 1390 1410 1430
AACACGCCcCTCCGGGTTCcCATGcTCTCGTcATGTCCATcTGGGAGcGACgtaagtaaccgcgcgacctccctagccacacaaagccgcatccggcgaggc
N N A L R V P M V L V M S I W D D

1450 1470 1490 1510 1530
acgccatcgctgctgctaacacgagaccgttcgtagCACTACgCCAACATGcTCTGGCTGAGcTCCATcTACCGCCCGAGAGAGGAGGcCCAGCCCGGCG
H Y A N M L W L D S I Y P P E K E G Q P G A

1550 1570 1590 1610 1630
CCGcCCGTGGGcGACTGCCCCACGgACTCGGGTGTCCCGCGGAGGTcGAGGcTcAGTTCcCCCGACGCGgtaagacttgcccccgaccaccaagcttccactt
A R G D C P T D S G V P A E V E A Q F P D A

1650 1670 1690 1710 1730
ctggatgccgaatgctaacacgcggaacagCCAGGTcGTcTGGTCCAAcATCGGcTTCGGGCCcCATCGGcTcCGAcCTcAGAcTcTAAAGCCGGTCCcATGC
Q V V W S N I R F G P I G S T Y D F *

1750 1770 1790 1810 1830
ACTGCAGCCcCTGGGCCCGTcACGCCCGcCCACcTCCcCTCGGGAAAcTcTCCGTcGTcGGGGcTCCAAAGcATTTTGGCCcTCAAGTTTTTTTCGTTC

FIG. 23C




FIG.24




FIG.25




FIG.26

10 30 50 70 90
CCATGGACGGAAGTGGGAAGTCTTCTGCCCGAGCTGAAGACCCAGAGCATCCAGACCAGGCAACCCAGTGCACCCAGGAGATGAAGGTCTACGAGAACAT
110 130 150 170 190
TGACGGCTGGCTGACAGCCTGCCCCGGAACGTCCCCATACCGGGTCCGAGCCGGCTCTGGTAAGTCAAGAGATGAATGCTACCTTCCACCT
210 230 250 270 290
TCCACCCAGCGCAATACTTTCTCCCTCCCGTGCCCGTATTCTTCAACGCCCCGAGACTGACAGACCCGCTCGTCCAGCGGCAACCCCGGCA
310 330 350 370 390
ACGGCGGGCAGCAACCCGGGCAACGGCGGGCGGGGGCTGCACCGTCCAGAAAGTGGGGCAGTGGCGGGCATCGGCTACTCGGGCTGCACCCACCTG
410 430 450 470 490
CAAGGCCGGCTCGACCTGCCTCGGCCCCAGAACGAGTACTACTCGCAGTGCCTGTAAAGCGGCGGTGGGCTAGGTGGCCGAGCGGGGGGTTCTTCATTGG
K A G S T C P A Q N E Y Y S Q C L *
510 530 550 570 590
TTGAGCAATAGAACAGGAITTCGGCTCGTTGGCAGCGGCGGCGCGGGGATGGTGTGTACAATTCAAGACCTCAGTACCGAGGGAACCTGGAAAGGA
610 630 650 670 690
GTCAGTCTGCTTGACGGAAGCTGGCTGCCCGTGGCGGGCGCTGGCAAGGTAGATAGCCCTTCATTGCTGTAAGTAGTATGCTATATACCTCTGCACATT
710 730 750 770 790
TGCAGCCCCATGGTGTGAACAACAAGTGAACAAGGCTTCCAGTTCAGCCCTCGGCAATTGTACAGATATCTTGGTCCATCTATATGTATGGGCATGAGC
810 830 850 870
GAGTCGAGAAATGTACCGGGAATAATCGTAGTGACCTGGGCACTGGCGGTTCTACCAACGTAGGATTGAAGTGAATCTCGAATTC

FIG. 27




FIG.28




FIG.29

1 2 3 4 5 6




FIG.30

FIG.31




FIG.32

CELLULASES, THE GENES ENCODING THEM AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of International Application No. PCT/FI96/00550, filed Oct. 17, 1996, and a continuation-in-part of U.S. application Ser. No. 08/732,181, filed Oct. 16, 1996, which claims the benefit of U.S. Provisional Application Nos. 60/005,335, filed Oct. 17, 1995; 60/007,926, filed Dec. 4, 1995; and 60/020,840, filed Jun. 28, 1996.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention is related to genes encoding novel neutral cellulases and compositions containing the novel neutral cellulases. These compositions are especially useful in the textile, detergent and pulp and paper industries.

2. Related Art

Cellulose is a linear polysaccharide of glucose residues connected by β -1,4 linkages. In nature, cellulose is usually associated with lignin together with hemicelluloses such as xylans and glucomannans. The practical use of cellulases has been hampered by the nature of the known cellulases, which are often mixtures of cellulases having a variety of activities and substrate specificities. For that reason, it is desirable to identify sources from which cellulases having only the desired activities may be obtained.

A wide variety of cellulases are known in the art, most of which are acid cellulases. However, some neutral and alkaline cellulases have also been identified. Celluzyme® is a commercially-available cellulase preparation from *Humicola insolens* (Novo Nordisk, A/S). GB 2,075,028 and EP 406,314 describe the use of a *Humicola insolens* cellulase as an enzymatic additive in a wash detergent to reduce the harshness (stiffness) of cotton-containing fabrics. The cloning of a cellulase containing endoglucanase activity from *Humicola insolens* is described in WO 93/11249 and EP 531,372. EP 510,091 describes a cellulase from *Bacillus* spp. NCIMB 40250 that is useful in detergent compositions. EP 220,016 describes cellulases that are useful as clarification agents for colored fabrics. WO 94/07998 describes modified cellulases that possess an improved alkaline activity. WO 95/02675 describes detergent compositions that contain two different cellulases: a first cellulase that is catalytically amenable to particulate soil removal, and a second cellulase that is catalytically amenable to color clarification. WO 92/18599 describes a detergent preparation that contains both a cellulase and a protease. Cellulases have also been used industrially as an aid for the removal of printing paste thickener and excess dye after textile printing (EP 576,526).

EP 383 828 describes granular detergent compositions, which contain surface-active agent, a fabric-softening clay material, and cellulase granulates containing calcium carbonate. U.S. Pat. No. 5,433,750 describes detergent compositions containing a surface active agent, a builder system, a softening clay, a clay flocculating agent and a high activity cellulase, preferably *Humicola insolens* cellulase. U.S. Pat. No. 5,520,838 describes granular detergent compositions, comprising surface-active agent, a builder and a cellulase, preferably a *Humicola insolens* cellulase, said compositions being in a compact form, having a relatively high density and containing a low amount of inorganic filler salt.

Cellulase enzymes are used in a wide variety of industries in addition to the textile industry. For example, cellulases are

used industrially for the deinking of newspapers and magazines (EP 521,999), for improving the drainage of pulp (WO 91/14822, WO 91/17243), and as a treatment for animal feed.

The unique properties of each cellulase make some more suitable for certain purposes than others. While the enzymes differ in a number of ways, one of the most important difference is pH optimum. Neutral cellulases have useful cellulase activity in the pH range 6–8, alkaline cellulases have useful cellulase activity in the pH range 7.5–10. Acid cellulases are active in the range of pH 4.5–6, but have little cellulase activity at higher pH values.

Neutral and acid cellulases are especially useful in the textile industry Klahorst, S. et al., *Textile Chemist and Colorist* 26:13–18, 1994; Nilsson, T. E., Aachen Textile Conference, *DWI Reports* 114:85–88 (1995); Videbæk, T. et al., *ITB Dyeing/Printing/Finishing*, January 1994, pp. 25–29; Klahorst, S. et al., *AATCC Int. Conf. & Exhibit*, Oct. 4–7, 1992, p. 243, Atlanta, Ga.; Kochavi, D. et al., *Am. Dyestuff Reporter*, September 1990, pp. 26–28; Tyndall, R. Michael, *Textile Chemist and Colorist* 24:23 (1992); Lange, N. K., in *Proc. Second TRICEL Symp. on Trichoderma reesei Cellulases and Other Hydrolases*, Espoo, Finland, 1993, ed. P. Suominen et al., Foundation for Biotechnical and Industrial Fermentation Research vol. 8, 1993, pp. 263–272. When used to treat fabric, cellulases attack the chains of cellulose molecules that form the cotton fibers, thereby affecting the characteristics of the fabric.

Traditionally, in “stonewashing,” pumice stones have been used to change the characteristics of the fabric. Gradually, cellulases are replacing pumice stones, which also give the fabric its desired final look but can cause damage to the machines, garments and sewage processing equipment. U.S. Pat. Nos. 4,832,864, 4,912,056, 5,006,126, 5,122,159, 5,213,581 and EP 307,564 disclose the use of cellulases in biostoning.

Cellulases are especially useful for stonewashing denim dyed with indigo as the dye mostly stays on the surface of the yarn and does not penetrate the fibers well. When used to treat cotton fabric, neutral cellulases generally require a longer wash time than do the acid cellulases. However, available neutral cellulases are less aggressive (active) against cotton than acid cellulases, and are reported not to compromise the strength of the fabric as readily as acid cellulases. Neutral cellulases have a broader pH profile and thus the pH increase that occurs during biostoning has little effect on the activity of the neutral enzyme.

The use of acid cellulases is hampered by their tendency to promote backstaining and a weakening of fabrics. In addition, the pH must be adjusted to to a range suitable for the function of the acid cellulases. Consequently, there is a clear demand for neutral cellulase enzyme preparations that do not cause backstaining or weakening of fabrics.

While it has become popular to use cellulases in the textile industry, simply changing the cellulase mixture that is used may produce a different finish. These problems have focused increasing attention on the search for reproducible mixtures of cellulases with desired properties. Thus there is a clear demand especially in the textile and detergent industry for novel cellulases active at neutral and alkaline pH values, not compromising the strength of fabrics, with good cleaning and/or fabric care and harshness reducing properties.

SUMMARY OF THE INVENTION

Recognizing the importance of identifying enzymes useful in textile biofinishing and biostoning and in detergent

applications, the inventors have screened fungal species for neutral and alkaline cellulases with enzymatic characteristics that would be useful in such technologies.

These studies have resulted in novel cellulases originating from the genera *Myceliophthora*, *Myriococcum*, *Melanocarpus*, *Sporotrichum* and *Chaetomium*.

The invention is further directed to the spent culture medium or enzyme preparations prepared from the native hosts producing such novel cellulases.

The invention is further directed to the use of such culture medium or the use of such enzyme preparations in the textile and detergent industry and in the pulp and paper industries.

These studies have further resulted in the identification of three novel cellulases that are especially useful in the textile and detergent industry. Purified preparations from *Melanocarpus* sp. or *Myriococcum* sp. have revealed a 20 kDa cellulase with endoglucanase activity (designated herein as "20K-cellulase"), a 50 kDa cellulase ("50K-cellulase"), and a second 50 kDa cellulase ("50K-cellulase B"). A novel gene product with high homology to the cellulase family, herein called "protein-with-CBD" (where CBD means "cellulose binding domain") was also discovered.

It is an object of the invention to provide enzyme preparations that contain one or more of the novel cellulases of the invention, especially the 20K-cellulase, the 50K-cellulase, the 50K-cellulase B and/or the protein-with-CBD.

It is a further object of this invention to provide a method for using such preparations for the finishing of textiles, especially the biostoning of denim, for the use said preparations in detergent compositions, and especially methods that use the 20K-cellulase, the 50K-cellulase, the 50K-cellulase B and/or the protein-with-CBD.

The invention is also directed to other neutral and/or alkaline cellulases having one or more of the amino acid sequences as described herein.

The invention is further directed to the genes encoding the 20K-cellulase, 50K-cellulase, 50K-cellulase B and the protein-with-CBD.

The invention is further directed to novel expression vectors comprising such genes and to novel hosts transformed with the vectors, especially hosts that are capable of high levels of expression of the proteins encoded by such genes.

The invention is further directed to the spent culture medium of such transformed hosts, the culture medium containing the novel 20K-cellulase, 50K-cellulase, the 50K-cellulase B and/or the protein-with-CBD, or enzyme compositions (enzyme preparations) containing one or more of these proteins that have been prepared from such culture media.

The invention is further directed to the use of such culture medium or the use of such enzyme preparations in the textile and detergent industry and in the pulp and paper industries.

BRIEF DESCRIPTION OF THE FIGURES

FIGS. 1(A and B) show the pH (FIG. 1A) and temperature (FIG. 1B) dependencies of the endoglucanase activities of ALKO4179, CBS 689.95.

FIGS. 2(A and B) show the pH (FIG. 2A) and temperature (FIG. 2B) dependencies of the endoglucanase activities of ALKO4124, CBS 687.95.

FIGS. 3(A and B) show the pH (FIG. 3A) and temperature (FIG. 3B) dependencies of the endoglucanase activities of ALKO4237, CBS 685.95.

FIGS. 4(A and B) show the pH (FIG. 4A) and temperature (FIG. 4B) dependencies of the endoglucanase activities of ALKO4265, CBS 730.95.

FIGS. 5(A and B) show the pH (FIG. 5A) and temperature (FIG. 5B) dependencies of the endoglucanase activities of ALKO4125, CBS 688.95.

FIGS. 6(A and B) show the wash effect and backstaining (FIG. 6A) and blueness (FIG. 6B) with the neutral cellulases.

FIGS. 7(A and B) show the wash effect and backstaining (FIG. 7A) and blueness (FIG. 7B) with Ecoston L with gradually increasing enzyme dosages. 1X corresponds to the enzyme dosage of the neutral cellulases in FIGS. 6A and 6B.

FIG. 8 shows the purification of 20K-cellulase from Peak II by chromatography on SP-Sepharose™. A sample containing 11.7 g of protein and 576,000 ECU was applied to a 4.5x31 cm column, which was developed as described in Example 9. Fractions of 15 ml were collected. Endoglucanase activities in the peak at fractions 148-161 are underestimated because crystallization occurred before the enzyme could be sufficiently diluted for assay. Crystalline material from these fractions contained 486,000 ECU.

FIGS. 9(A and B) show SDS-PAGE analysis of the 20K-cellulase. The molecular masses of the standards are shown in kDa.

A Partially crystalline material precipitated from the active S-Sepharose™ fractions (lane 1).

B Fractions from chromatography of the partially crystalline material on G50 Sephadex. Fractions shown in lanes 19 and 25 contained no endoglucanase activity. For the other fractions, the amounts of activity (in ECU) applied to the gel was as follows: fraction 27, 0.4; 29, 2.4 (as 3.0 µg of protein); 30, 2.1; 31, 1.9; 33, 0.46; and 35, 1.1.

FIG. 10 shows the separation of 50K-cellulase and 50K-cellulase B from Peak III/IV by chromatography on SP-Sepharose™. A sample containing 200 mg of protein and 14,800 ECU was applied to the 2.5x11 cm column, which was developed as described in Example 9. Fractions of 6.8 ml were collected. A minor amount of 50K-cellulase eluted before the NaCl gradient, whereas most of the 50K-cellulase eluted at about 50 mM NaCl. 50K-cellulase B was found in the major protein peak at about 80 mM NaCl.

FIG. 11 shows an SDS-PAGE analysis of purified 50K-cellulase (11A) and 50K-cellulase B (11B). Lane numbers indicate the fractions (3.3 ml) eluted from Phenyl-Sepharose. For fractions 36-41, 2.5 µl of each fraction was applied to the gel. For the other fractions, 2 µl was applied. The 50K-cellulase peak was found in fractions 37-38 (11A) (containing 780 and 880 ECU/ml, respectively). The 50K-cellulase B peak was in fractions 30 and 31 (11B), which contained negligible activity (less than 4 ECU/ml).

FIG. 12 shows the temperature dependence of the endoglucanase activity of 50K-cellulase at pH 7.0 and a reaction time of 60 min.

FIG. 13 shows the pH dependence of the endoglucanase activity of 50K-cellulase at 50° C. (♦) and 70° C. (□).

FIG. 14 shows a Western analysis using 20K-cellulase antiserum as a probe. Lanes 1, 2 and 3 contain 25 µg of protein from the DEAE-Sepharose peaks I, III and IV, respectively. Lanes 4 and 5 contain 2.0 and 0.2 µg of pure 50K-cellulase and lane 6 contains 0.6 µg of pure 50K-cellulase B. Lanes 7 and 8 contain about 25 µg protein from the whole growth medium of ALKO4237 and ALKO4124, respectively.

FIG. 15 shows the temperature dependence of the endoglucanase activity of 20K-cellulase at pH 7 (10 min reaction times).

FIGS. 16(A and B) show the pH-dependence of the endoglucanase activity of the 20K-cellulase for the reaction time of (a) 10 minutes or (b) 60 minutes.

FIG. 17 shows amino acid sequence data derived from sequencing the 20K-cellulase described in the exemplary material herein. Sequence 429 (SEQ ID NO:1) is from the N terminus of the protein and the other sequences are from internal tryptic peptides. Sequence #430 corresponds to SEQ ID NO: 2; sequence #431 corresponds to SEQ ID NO: 3; sequence #432 corresponds to SEQ ID NO: 4; sequence #433 corresponds to SEQ ID NO: 5; sequence #439 corresponds to SEQ ID NO: 6; fr 9 corresponds to SEQ ID NO: 7; fr 14 corresponds to SEQ ID NO: 8; fr 16 corresponds to SEQ ID NO: 9; fr 17 corresponds to SEQ ID NO: 10; fr 28 corresponds to SEQ ID NO: 11 and fr 30 corresponds to SEQ ID NO: 12.

FIG. 18 shows the restriction maps of the *Melanocarpus albomyces* DNA in plasmids pALK1221, pALK1222 and pALK1223, which carry the 20K-cellulase gene.

FIG. 19 shows the DNA sequence of the 20K-cellulase gene (SEQ ID NO:30). The arrow indicates the predicted signal peptidase processing site.

FIG. 20 shows the restriction maps of the *Melanocarpus albomyces* DNA in plasmids pALK1233, pALK1234, pALK1226 and pALK1227, which carry the 50K-cellulase gene.

FIGS. 21 (A, B and C) show the DNA sequence of the 50K-cellulase gene. The arrow indicates the predicted signal peptidase processing site.

FIG. 22 shows the restriction maps of the *Melanocarpus albomyces* DNA in plasmids pALK1229 and pALK1236, which carry the 50K-cellulase B gene.

FIGS. 23 (A, B and C) show the DNA sequence of the 50K-cellulase B gene. The arrow indicates the predicted signal peptide processing site.

FIG. 24 shows the plasmid map of pTTC01.

FIG. 25 shows the plasmid map of pMS2.

FIG. 26 shows the restriction map of the *Melanocarpus albomyces* DNA in plasmid pALK1230, which carries DNA encoding the protein-with-CBD. The sequence presented in FIG. 27 (SEQ ID NO: 36) is marked with an arrow in FIG. 26.

FIG. 27 shows the DNA sequence of the the protein-with-CBD cellulase gene (SEQ ID NO:36) in pALK1230.

FIG. 28 shows the plasmid map of pALK1231.

FIG. 29 shows the plasmid map of pALK1235.

FIG. 30 shows a Western analysis using 20K-cellulase antiserum as a probe. Lanes 1 and 2 contain about 10 μ g protein from the whole growth medium of transformants ALKO3620/pALK1235/49 and ALKO3620/pALK1235/40. Lane 3 contains about 10 μ g protein from the whole growth medium of ALKO3620. Lanes 4 and 5 contain about 10 μ g protein from the whole growth medium of transformants ALKO3620/pALK1231/16 and ALKO3620/pALK1231/14. Lane 6 contains 100 ng of pure 20K-cellulase.

FIG. 31 shows the plasmid map of pALK1238.

FIG. 32 shows the plasmid map of pALK1240.

DEPOSITS

ALKO4179, *Myceliophthora thermophila* was deposited as CBS 689.95 on Oct. 12, 1995, at the Centraalbureau voor Schimmelcultures, P.O. Box 273, 3740 AG BAARN.

ALKO4124, *Myriococcum* sp. was deposited as CBS 687.95 on Oct. 12, 1995, at the Centraalbureau voor Schimmelcultures, P.O. Box 273, 3740 AG BAARN.

ALKO4237, *Melanocarpus albomyces* (= *Myriococcum albomyces*=*Thielavia albomyces*; Guarro et al., 1996,

Mycol. Res. 100(1):75.) was deposited as CBS 685.95 on Oct. 11, 1995, at the Centraalbureau voor Schimmelcultures, P.O. Box 273, 3740 AG BAARN.

ALKO4125, *Sporotrichum thermophile* was deposited as CBS 688.95 on Oct. 12, 1995, at the Centraalbureau voor Schimmelcultures, P.O. Box 273, 3740 AG BAARN.

ALKO4265, *Chaetomium thermophilum* La Touche was deposited as CBS 730.95 on Nov. 8, 1995, at the Centraalbureau voor Schimmelcultures, P.O. Box 273, 3740 AG BAARN.

Plasmid pALK1221 was deposited as DSM 11024 on Jun. 21, 1996 and λ 4237/5.1 was deposited as DSM 11012 on Jun. 21, 1996, at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1B, D-38124 Braunschweig, Germany. Both contain the 20K-cellulase gene from *Melanocarpus albomyces* CBS 685.95.

Plasmid pALK1227 was deposited as DSM 11025 on Jun. 21, 1996 and λ 4237/35 was deposited as DSM 11014 on Jun. 21, 1996, at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1B, D-38124 Braunschweig, Germany. Both contain the 50K-cellulase gene from *Melanocarpus albomyces* CBS 685.95.

Plasmid pALK1229 was deposited as DSM 11026 on Jun. 21, 1996 and λ 4237/3 was deposited as DSM 11011 on Jun. 21, 1996, and λ 4237/118 was deposited as DSM 11013 on Jun. 21, 1996, at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1B, D-38124 Braunschweig, Germany. pALK1229 contains DNA coding for the 50K-cellulase B, λ 14237/3 and λ 4237/18 contain the 50K-cellulase B gene from *Melanocarpus albomyces* CBS 685.95.

Plasmid pALK1230 was deposited as DSM 11027 on Jun. 21, 1996 at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1B, D-38124 Braunschweig, Germany. pALK1230 contains the protein-with-CBD gene from *Melanocarpus albomyces* CBS 685.95.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In the description that follows, a number of terms used in textile industry technology are extensively utilized. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

Biostoning. "Biostoning" of fabric or garment means the use of enzymes in place of, or in addition to, the use of pumice stones for the treatment of fabric or garment, especially denim.

Biofinishing. "Biofinishing" refers to the use of enzymes in a controlled hydrolysis of cellulosic fibers in order to modify the fabric or yarn surface in a manner that prevents permanently pilling, improves fabric handle like softness and smoothness, clears the surface structure by reducing fuzzing, which results in clarification of colours, improves the drapability of the fabric, improves moisture absorbability and which may improve also the dyeability.

Backstaining. Released dye has a tendency to redeposit on the surface of the fabric fibers. This effect is termed "backstaining."

Detergent. By "detergent" is meant a cleansing agent that can contain surface active agents (anionic, non-ionic, cationic and ampholytic surfactants), builders and other optional ingredients such as antiredeposition and soil suspension agents, optical brighteners, bleaching agents, dyes

and pigments and hydrolases. Suitable listing of the contents of detergents is given in U.S. Pat. No. 5,433,750, a suitable list of surfactants is given in U.S. Pat. No. 3,664,961.

Enzyme preparation. By "enzyme preparation" is meant a composition containing enzymes. Preferably, the enzymes have been extracted from (either partially or completely purified from) a microbe or the medium used to grow such microbe. "Extracted from" means that the desired enzymes are separated from the cellular mass. This can be performed by any method that achieves this goal, including breaking cells and also simply removing the culture medium from spent cells. Therefore, the term "enzyme preparation" includes compositions containing medium previously used to culture a desired microbe(s) and any enzymes that have been released from the microbial cells into such medium during the culture or downstream processing steps.

By a host that is "substantially incapable" of synthesizing one or more enzymes is meant a host in which the activity of one or more of the listed enzymes is depressed, deficient, or absent when compared to the wild-type.

By an amino acid sequence that is an "equivalent" of a specific amino acid sequence is meant an amino acid sequence that is not identical to the specific amino acid sequence, but rather contains at least some amino acid changes (deletions, substitutions, inversions, insertions, etc) that do not essentially affect the biological activity of the protein as compared to a similar activity of the specific amino acid sequence, when used for a desired purpose. The biological activity of a cellulase, is its catalytic activity, and/or its ability to bind to cellulosic material. The biological activity of the 50K-cellulase B further includes its ability to act synergistically with the cellulases. Preferably, an "equivalent" amino acid sequence contains at least 80%-99% identity at the amino acid level to the specific amino acid sequence, most preferably at least 90% and in an especially highly preferable embodiment, at least 95% identity, at the amino acid level.

Cloning vehicle. A cloning vehicle is a plasmid or phage DNA or other DNA sequence (such as a linear DNA) that provides an appropriate nucleic acid carrier environment for the transfer of a gene of interest into a host cell. The cloning vehicles of the invention may be designed to replicate autonomously in prokaryotic and eukaryotic hosts. In fungal hosts such as *Trichoderma*, the cloning vehicles generally do not autonomously replicate and instead, merely provide a vehicle for the transport of the gene of interest into the *Trichoderma* host for subsequent insertion into the *Trichoderma* genome. The cloning vehicle may be further characterized by one or a small number of endonuclease recognition sites at which such DNA sequences may be cut in a determinable fashion without loss of an essential biological function of the vehicle, and into which DNA may be spliced in order to bring about replication and cloning of such DNA. The cloning vehicle may further contain a marker suitable for use in the identification of cells transformed with the cloning vehicle. Markers, for example, are antibiotic resistance. Alternatively, such markers may be provided on a cloning vehicle which is separate from that supplying the gene of interest. The word "vector" is sometimes used for "cloning vehicle."

Expression vehicle. An expression vehicle is a cloning vehicle or vector similar to a cloning vehicle but which is capable of expressing a gene of interest, after transformation into a desired host. When a fungal host is used, the gene of interest is preferably provided to a fungal host as part of a cloning or expression vehicle that integrates into the fungal

chromosome, or allows the gene of interest to integrate into the host chromosome. Sequences that are part of the cloning vehicle or expression vehicle may also be integrated with the gene of interest during the integration process. In *T. reesei*, sites of integration to which the gene of interest can be directed include the *cbh* and/or the *egl* loci. Most preferably, the gene of interest is directed to replace one or more genes encoding undesirable characteristics.

The gene of interest is also preferably placed under the control of (i.e., operably linked to) certain control sequences such as promoter sequences provided by the vector (which integrate with the gene of interest). Alternatively, the control sequences can be those at the insertion site.

The expression control sequences of an expression vector will vary depending on whether the vector is designed to express a certain gene in a prokaryotic or in a eukaryotic host (for example, a shuttle vector may provide a gene for selection in bacterial hosts). Expression control sequences can contain transcriptional regulatory elements such as, promoters, enhancer elements, and transcriptional termination sequences, and/or translational regulatory elements, such as, for example, translational initiation and termination sites.

According to the invention, there are provided neutral and alkaline cellulases, and methods for producing such useful neutral and alkaline cellulases, that are desirable for the treatment of textile materials.

The native hosts that produce the proteins of the invention are:

- 1) ALKO4179, *Myceliophthora thermophila*; deposited as CBS 689.95 at the Centraalbureau voor Schimmelcultures, P.O. Box 273, 3740 AG BAARN.
- 2) ALKO4124, *Myriococcum* sp.; deposited as CBS 687.95;
- 3) ALKO4237, *Melanocarpus albomyces*, deposited as CBS 685.95;
- 4) ALKO4125, *Sporotrichum thermophila*, deposited as CBS 688.95; and
- 5) ALKO4265, *Chaetomium thermophilum* La Touche, deposited as CBS 730.95

One specific preferred embodiment of the invention is the spent culture medium of the native hosts or enzyme preparations prepared from the culture medium.

In specific preferred embodiments of the invention, the purified 20K-cellulase, 50K-cellulase, 50K-cellulase B and/or protein-with-CBD is provided. These proteins can be obtained for example from *Melanocarpus* sp. or *Myriococcum* sp. as described herein, and especially in Example 9.

Amino acid sequence data have been generated from the cellulases described herein. Accordingly, the invention is also directed to neutral or alkaline cellulases containing one or more of the amino acid sequences shown herein. Thus, the invention is intended to be directed to any neutral or alkaline cellulase that is a functional equivalent of the 20K-cellulase, the 50K-cellulase, the 50K-cellulase B and/or protein-with-CBD and having one or more of the amino acid sequences described herein, or substantially the same sequence. Such neutral or alkaline cellulases can be derived from other strains of the same species or from divergent organisms.

In further preferred embodiments, the 20K-cellulase is provided with the material from separate peaks formed during the exemplified purification procedures (e.g., DEAE-Sephacrose Pools I, III, or IV in Table VIII herein). In still further embodiments, other proteins in the *Melanocarpus albomyces* ALKO 4237 medium may be used, either alone or in combination with other such proteins.

In further preferred embodiments, the 50K-cellulase is provided with the material from separate peaks formed during the exemplified purification procedures. In still further embodiments, other proteins in the ALKO 4237 medium may be used, either alone or in combination with other such proteins.

In further preferred embodiments, the 50K-cellulase B is provided with the material from separate peaks formed during the exemplified purification procedures. In still further embodiments, other proteins in the ALKO 4237 medium may be used, either alone or in combination with other such proteins.

As described herein, ALKO 4265, *Chaetomium thermophilum* La Touche, deposited as CBS 730.95, is used herein as an example of a neutral cellulase that is not preferred in biostoning method of the invention because it causes back-staining. However, there is evidence that it is useful in other applications (e.g. in detergents).

The process for genetically engineering the hosts of the invention is facilitated through the cloning of genetic sequences that encode the desired protein and through the expression of such genetic sequences. As used herein the term "genetic sequences" is intended to refer to a nucleic acid molecule (preferably DNA). Genetic sequences that encode the desired protein are derived from a variety of sources. These sources include genomic DNA, cDNA, synthetic DNA and combinations thereof. Vector systems may be used to produce hosts for the production of the enzyme preparations of the invention. Such vector construction (a) may further provide a separate vector construction (b) which encodes at least one desired gene to be integrated to the genome of the host and (c) a selectable marker coupled to (a) or (b). Alternatively, a separate vector may be used for the marker.

A nucleic acid molecule, such as DNA, is said to be "capable of expressing" a polypeptide if it contains expression control sequences which contain transcriptional regulatory information and such sequences are "operably linked" to the nucleotide sequence which encodes the polypeptide.

An operable linkage is a linkage in which a sequence is connected to a regulatory sequence (or sequences) in such a way as to place expression of the sequence under the influence or control of the regulatory sequence. Two DNA sequences (such as a protein encoding sequence and a promoter region sequence linked to the 5' end of the encoding sequence) are said to be operably linked if induction of promoter function results in the transcription of the protein encoding sequence mRNA and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the expression regulatory sequences to direct the expression of the mRNA, antisense RNA, or protein, or (3) interfere with the ability of the template to be transcribed by the promoter region sequence. Thus, a promoter region would be operably linked to a DNA sequence if the promoter were capable of effecting transcription of that DNA sequence.

The precise nature of the regulatory regions needed for gene expression may vary between species or cell types, but shall in general include, as necessary, 5' non-transcribing and 5' non-translating (non-coding) sequences involved with initiation of transcription and translation respectively. Expression of the protein in the transformed hosts requires the use of regulatory regions functional in such hosts. A wide variety of transcriptional and translational regulatory sequences can be employed. In eukaryotes, where transcription is not linked to translation, such control regions may or

may not provide an initiator methionine (AUG) codon, depending on whether the cloned sequence contains such a methionine. Such regions will, in general, include a promoter region sufficient to direct the initiation of RNA synthesis in the host cell.

As is widely known, translation of eukaryotic mRNA is initiated at the codon which encodes the first methionine. For this reason, it is preferable to ensure that the linkage between a eukaryotic promoter and a DNA sequence which encodes the protein, or a functional derivative thereof, does not contain any intervening codons which are capable of encoding a methionine. The presence of such codons results either in a formation of a fusion protein (if the AUG codon is in the same reading frame as the protein encoding DNA sequence) or a frame-shift mutation (if the AUG codon is not in the same reading frame as the protein encoding sequence).

In a preferred embodiment, a desired protein is secreted into the surrounding medium due to the presence of a secretion signal sequence. If a desired protein does not possess its own signal sequence, or if such signal sequence does not function well in the host, then the protein's coding sequence may be operably linked to a signal sequence homologous or heterologous to the host. The desired coding sequence may be linked to any signal sequence which will allow secretion of the protein from the host. Such signal sequences may be designed with or without specific protease sites such that the signal peptide sequence is amenable to subsequent removal. Alternatively, a host that leaks the protein into the medium may be used, for example a host with a mutation in its membrane.

If desired, the non-transcribed and/or non-translated regions 3' to the sequence coding for a protein can be obtained by the above-described cloning methods. The 3'-non-transcribed region may be retained for its transcriptional termination regulatory sequence elements; the 3'-non-translated region may be retained for its translational termination regulatory sequence elements, or for those elements which direct polyadenylation in eukaryotic cells.

The vectors of the invention may further comprise other operably linked regulatory elements such as enhancer sequences.

In a preferred embodiment, genetically stable transformants are constructed whereby a desired protein's DNA is integrated into the host chromosome. The coding sequence for the desired protein may be from any source. Such integration may occur de novo within the cell or, in a most preferred embodiment, be assisted by transformation with a vector which functionally inserts itself into the host chromosome, for example, DNA elements which promote integration of DNA sequences in chromosomes.

Cells that have stably integrated the introduced DNA into their chromosomes are selected by also introducing one or more markers which allow for selection of host cells which contain the expression vector in the chromosome, for example the marker may provide biocide resistance, e.g., resistance to antibiotics, or heavy metals, such as copper, or the like. The selectable marker gene can either be directly linked to the DNA gene sequences to be expressed, or introduced into the same cell by co-transformation.

Factors of importance in selecting a particular plasmid or viral vector include: the ease with which recipient cells that contain the vector may be recognized and selected from those recipient cells which do not contain the vector; the number of copies of the vector which are desired in a particular host; and whether it is desirable to be able to "shuttle" the vector between host cells of different species.

Once the vector or DNA sequence containing the construct(s) is prepared for expression, the DNA construct

(s) is introduced into an appropriate host cell by any of a variety of suitable means, including transformation as described above. After the introduction of the vector, recipient cells are grown in a selective medium, which selects for the growth of transformed cells. Expression of the cloned gene sequence(s) results in the production of the desired protein, or in the production of a fragment of this protein. This expression can take place in a continuous manner in the transformed cells, or in a controlled manner.

Accordingly, the protein encoding sequences described herein may be operably linked to any desired vector and transformed into a selected host, so as to provide for expression of such proteins in that host.

The subject matter of the invention are also nucleic acid molecules coding for proteins having the biological activity of a cellulase and that hybridize to any of the nucleic acid molecules described above or which are defined in the following:

A nucleic acid molecule encoding a polypeptide having the enzymatic activity of a cellulase, selected from the group consisting of:

- (a) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence as depicted in FIG. 19 (SEQ ID NO:31) or 21 (SEQ ID NO:33);
- (b) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence as depicted in FIG. 23 (SEQ ID NO:35) or 27 (SEQ ID NO:37);
- (c) nucleic acid molecules comprising the coding sequence of the nucleotide sequence as depicted in FIG. 19 (SEQ ID NO:30) or 21 (SEQ ID NO:32);
- (d) nucleic acid molecules comprising the coding sequence of the nucleotide sequence as depicted in FIG. 23 (SEQ ID NO:34) or 27 (SEQ ID NO:36);
- (e) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence encoded by the DNA insert contained in DSM 11024, DSM 11012, DSM 11025 or DSM 11014;
- (f) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence encoded by the DNA insert contained in DSM 11026, DSM 11011, DSM 11013 or DSM 11027;
- (g) nucleic acid molecules comprising the coding sequence of the DNA insert contained in DSM 11024, DSM 11012, DSM 11025 or DSM 11014;
- (h) nucleic acid molecules comprising the coding sequence of the DNA insert contained in DSM 11026, DSM 11011, DSM 11013 or DSM 11027;
- (i) nucleic acid molecules hybridizing to a molecule of any one of (a), (c), (e) or (g); and
- (j) nucleic acid molecules the coding sequence of which differs from the coding sequence of a nucleic acid molecule of any one of (a) to
- (i) due to the degeneracy of the genetic code.
- (k) nucleic acid molecules encoding a polypeptide having cellulase activity and having an amino acid sequence which shows at least 80% identity to a sequence as depicted in FIG. 19 (SEQ ID NO:31), 21 (SEQ ID NO:33), 23 (SEQ ID NO:35) or 27 (SEQ ID NO:37).

The term "hybridization" in this context means hybridization under conventional hybridization conditions, preferably under stringent conditions such as described by, e.g. Sambrook et al (1989, *Molecular Cloning, A Laboratory Manual* 2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). These nucleic acid molecules that hybridize to the nucleic acid molecules according to the

present invention in principle can be derived from any organism possessing such nucleic acid molecules. Preferably, they are derived from fungi, namely from those of the genera *Melanocarpus*, *Myriococcum*, *Sporotrichum*, *Myceliophthora* and *Chaetomium*. Nucleic acid molecules hybridizing to the nucleic acid molecules of the present invention can be isolated, e.g., from genomic libraries or cDNA libraries of various organisms, namely fungi.

Such nucleic acid molecules can be identified and isolated by using the nucleic acid molecules of the present invention or fragments of these molecules or the reverse complements of these molecules, e.g. by hybridization according to standard techniques (see Sambrook et al.(1989)).

As hybridization probe, e.g. nucleic acid molecules can be used that have exactly or substantially the same nucleotide sequence indicated in the Figures or fragments of said sequence. The fragments used as hybridization probes can also be synthetic fragments obtained by conventional synthesis techniques and the sequence of which is substantially identical to that of the nucleic acid molecules according to the invention. Once genes hybridizing to the nucleic acid molecules of the invention have been identified and isolated it is necessary to determine the sequence and to analyze the properties of the proteins coded for by said sequence.

The term "hybridizing DNA molecule" includes fragments, derivatives and allelic variants of the above-described nucleic acid molecules that code for the above-described protein or a biologically active fragment thereof. Fragments are understood to be parts of nucleic acid molecules long enough to code for the described protein or a biologically active fragment thereof. The term "derivative" means in this context that the nucleotide sequences of these molecules differ from the sequences of the above-described nucleic acid molecules in one or more positions and are highly homologous to said sequence. Homology is understood to refer to a sequence identity of at least 40%, particularly an identity of at least 60%, preferably more than 80% and still more preferably more than 90%. The deviations from the nucleic acid molecules described above can be the result of deletion, substitution, insertion, addition or combination.

Homology furthermore means that the respective nucleotide sequences or encoded proteins are functionally and/or structurally equivalent. The nucleic acid molecules that are homologous to the nucleic acid molecules described above and that are derivatives of said nucleic acid molecules are regularly variations of said molecules which represent modifications having the same biological function. They may be naturally occurring variations, such as sequences of other organisms or mutations. These mutations may occur naturally or may be achieved by specific mutagenesis. Furthermore, these variations may be synthetically produced sequences. The allelic variants may be naturally occurring variants as well as synthetically produced or genetically engineered variants.

The proteins encoded by the various variants of the nucleic acid molecules of the invention share specific common characteristics, such as enzymatic activity, molecular weight, immunological reactivity, conformation, etc., as well as physical properties, such as electrophoretic mobility, chromatographic behaviour, sedimentation coefficients, solubility, spectroscopic properties, stability, pH optimum, temperature optimum, etc. Enzymatic activity of the cellulase can be detected e.g. as described on page 11 and in Examples 1 and 25.

The present invention furthermore relates to nucleic acid molecules the sequences of which differ from the sequences

of the above-identified molecules due to degeneracy of the genetic code, and which code for a protein having the biological activity of a cellulase.

The nucleic acid molecules of the invention are preferably RNA or DNA molecules, most preferably genomic DNA or cDNA.

The present invention also relates to antibodies which specifically recognize one of the above-described proteins according to the invention as well as to antibody fragments which have this property. These antibodies may be monoclonal or polyclonal. Methods for their production are well known in the art and are described in detail, for example, in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor Laboratory (1988).

Furthermore, the present invention relates to oligonucleotides which specifically hybridize with a nucleic acid molecule according to the invention or with the complementary strand of such a nucleic acid molecule. In this respect the term "specifically hybridize" means that such an oligonucleotide hybridizes under stringent hybridization conditions specifically to a nucleic acid molecule of the invention and does not show under such conditions cross-hybridization with sequences coding for other polypeptides. Preferably such oligonucleotides have a length of at least 10 nucleotides, more preferably of at least 15 nucleotides and most preferably of at least 30 nucleotides. They are preferably no longer than 100 nucleotides, more preferably no longer than 80 nucleotides and most preferably no longer than 60 nucleotides. In order to ensure that they specifically hybridize to a nucleic acid molecule of the present invention such oligonucleotides show over their total length an identity of at least 80%, preferably of at least 95% and most preferably of at least 99% with a corresponding nucleotide sequence of a nucleic acid molecule of the present invention. These oligonucleotides may be used, e.g., as probes for screening for sequences encoding cellulases in genomic or cDNA libraries or as PCR primers.

The protein encoding sequences described herein may be fused in frame to other sequences so as to construct DNA encoding a fusion protein. For example, a recombinant vector encoding a 50K-cellulase, a 20K-cellulase, a 50K-cellulase B or the protein-with-CBD gene can be prepared as above, except that the protein encoding sequence is fused with the sequence of a *T. reesei* cellulase, hemicellulase or mannanase, or at least one functional domain of such cellulase, hemicellulase, or mannanase as described in U.S. Pat. No. 5,298,405, WO 93/24622 and in GenBank submission L25310, each incorporated herein by reference. Especially, the cellulase, hemicellulase, or mannanase is selected from the group consisting of CBHI, CBHII, EGI, EGII, XYLI, XYLII and MANI, or a domain thereof, such as the secretion signal or the core sequence. Mannanase has the same domain structure as that of the cellulases: a core domain, containing the active site, a hinge domain containing a serine-threonine rich region, and a tail, containing the binding domain.

Fusion peptides can be constructed that contain a mannanase or cellobiohydrolase or endoglucanase or xylanase core domain or the core and the hinge domains from the same, fused to the desired protein encoding sequence of the invention. The result is a protein that contains mannanase or cellobiohydrolase or endoglucanase or xylanase core or core and hinge regions, and a 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD sequence. The fusion protein contains both the mannanase or cellobiohydrolase or endoglucanase or xylanase, and the 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD activities of the various domains as provided in the fusion construct.

Fusion proteins can also be constructed such that the mannanase or cellobiohydrolase or endoglucanase or xylanase tail or a desired fragment thereof, is included, placed before the 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD sequence, especially so as to allow use of a nonspecific protease site in the tail as a protease site for the recovery of the 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD sequence from the expressed fusion protein. Alternatively, fusion proteins can be constructed that provide for a protease site in a linker that is placed before the 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD sequence, with or without tail sequences.

New properties for the 20K- and 50K-cellulases and for the 50K-cellulase B can be created by fusing domains, such as a cellulose binding domain (CBD), preferably with its linker, to the proteins of the invention. Preferably, such CBD's and linkers are the corresponding CBD and linker domains of a *Trichoderma* cellulase, mannanase or of the *Melanocarpus albomyces* protein-with-CBD.

The invention provides methods for producing enzyme preparations that are partially or completely deficient in an undesirable cellulolytic activity (that is, in the ability to degrade cellulose) and enriched in the 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD protein, as desired for the textile or detergent industry or for pulp and paper processing. By "deficient in cellulolytic activity" is meant a reduced, lowered, or repressed capacity to degrade cellulose to smaller oligosaccharides. Such cellulolytic activity deficient preparations, and the making of same by recombinant DNA methods, are described in U.S. Pat. No. 5,298,405, incorporated herein by reference. Preferably, the preparation is deficient in EG activities, and/or CBHI activity.

As described herein, the 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD may be provided directly by the hosts of the invention. Alternatively, spent medium from the growth of the hosts, or purified 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD therefrom, can be used. Further, if desired activities are present in more than one recombinant host, such preparations may be isolated from the appropriate hosts and combined prior to use in the method of the invention.

To obtain the enzyme preparations of the invention, the native or recombinant hosts described above having the desired properties (that is, hosts capable of expressing economically feasible quantities of the desired 50K-cellulase, 20K-cellulase, 50K-cellulase B or protein-with-CBD, and optionally, those that are substantially incapable of expressing one or more other, undesired cellulase enzymes) are cultivated under suitable conditions, the desired enzymes are secreted from the hosts into the culture medium, and the enzyme preparation is recovered from said culture medium by methods known in the art.

The enzyme preparations of the invention can be produced by cultivating the recombinant hosts or native strains in a fermentor on a suitable growth medium (such as, for example, shown in Example 1 or in Example 30).

The enzyme preparation can be the culture medium with or without the native or transformed host cells, or is recovered from the same by the application of methods well known in the art. However, because the 50K-cellulase, 20K-cellulase or 50K-cellulase B are secreted into the culture media and display activity in the ambient conditions of the cellulolytic liquor, it is an advantage of the invention that the enzyme preparations of the invention may be utilized directly from the culture medium with no further

purification. If desired, such preparations may be lyophilized or the enzymatic activity otherwise concentrated and/or stabilized for storage. The enzyme preparations of the invention are very economical to provide and use because (1) the enzymes may be used in a crude form; isolation of a specific enzyme from the culture medium is unnecessary and (2) because the enzymes are secreted into the culture medium, only the culture medium need be recovered to obtain the desired enzyme preparation; there is no need to extract an enzyme from the hosts. Preferably the host for such production is *Trichoderma*, and especially *T. reesei*.

The enzyme preparations of the invention may be provided as a liquid or as a solid, for example, in a dried powder or granular or liquid form, especially nondusting granules, or a stabilized liquid, or the enzyme preparation may be otherwise concentrated or stabilized for storage or use. It is envisioned that enzyme preparations containing one or more of the neutral cellulases of the invention can be further enriched or made partially or completely deficient in specific enzymatic activities, so as to satisfy the requirements of a specific utility in various applications e.g. in the textile industry. A mixture of enzyme activities secreted by a host and especially a fungus, can be chosen to be advantageous in a particular industrial application, for example biostoning.

The enzyme preparations of the invention can be adjusted to satisfy the requirements of specific needs in various applications in the textile, detergent or the pulp and paper industry.

Blends may be prepared with other macromolecules that are not all secreted from the same host (for example, other enzymes such as endoglucanases, proteases, lipases, peroxidases, oxidases or amylases) or chemicals that may enhance the performance, stability, or buffering of the desired enzyme preparation. Non-dusting granules may be coated. Liquid enzyme preparations can be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid, according to established methods. Liquid detergents generally contain up to 90% water and 0-20% organic solvent. Protected forms of the enzymes of the invention may be prepared as described in EP 238,216.

The enzyme preparations of the invention can contain a surfactant which can be anionic, non-ionic, cationic, amphoteric or a mixture of these types, especially when used as a detergent composition. Useful detergent compositions are described e.g. in WO 94/07998, U.S. Pat. Nos. 5,443,750 and 3,664,961.

If required, a desired enzyme may be further purified in accordance with conventional conditions, such as extraction, precipitation, chromatography, affinity chromatography, electrophoresis, or the like.

The enzyme preparations of this invention are especially useful in textile industry preferably in biostoning; and in biofinishing or in detergent industry. Other useful areas are in pulp and paper industry.

Non-enzymatic stonewashing has three steps: desizing, abrasion and aftertreatment. The first step, desizing, involves the removal of the starch coating, or that of its derivatives, by amylase. The second step, abrasion, when performed without cellulase, is generally performed by washing the denim with pumice stones, and, when lightening is desired, bleach. The abrasive effect is the result not only of the effect of the stones but also the rubbing together of the denim fabric. Abrasion is generally followed by the third step, a washing step to remove excess dye, during which softeners or optical brighteners can be added.

In enzymatic stonewashing, or biostoning, abrasion with pumice stones is completely or partially eliminated and

cellulase is added to facilitate the abrasion of indigo dye from the fiber surface. After this treatment, the cellulase is removed with a detergent wash to ensure that the mechanical strength of the fiber is not further compromised by the continued presence of the enzyme. Treatment with a cellulase(s) can completely replace treatment with pumice stones (for example, 1 kg commercial enzyme per 100 kg stones). However, cellulase treatment can be combined with pumice stone treatment when it is desired to produce a heavily abraded finish. A peach skin effect in which a fine protruding hair-like covering is created is also achieved by a wash combining a neutral cellulase with pumice stones. The cellulases of this invention are useful especially to minimize backstaining and enhance lightening (abrasion) in biostoning.

Biostoning is preferably performed from about pH 4.5-9.5, and most preferably between pH 6.0-8.5. The temperature of the reaction can range from about 40-80° C., preferably between 50-70° C., and most preferably between 50-60° C. The liquid ratio (the ratio of the volume of liquid per weight of fabric) may range from about 2:1-20:1, preferably 4:1-10:1, and most preferably 4:1-7:1. The enzyme dosage can range from about 25-1500 nkat/g fabric, preferably 50-500 nkat/g fabric and most preferably 75-300 nkat/g fabric.

The cellulases of the invention are useful in the textile industry for biofinishing of fabrics or garments e.g. depilling, defuzzing, colour clarification, harshness reduction, the creation of different finishes (for example, a 'peach skin,' 'worn out,' 'sand washed,' or 'antique look' effect) and biofinishing of yarn (for example reduction of hairiness, improvement of smoothness). The cellulases of this invention can be used in biofinishing in acidic and in neutral conditions.

The cellulases of this invention are useful in detergent compositions to improve the textile cleaning effect e.g. soil removal, to improve the fabric-care properties by reducing the harshness of the textiles, the cellulases having also defuzzing and colour clarification and restoring effects.

The textile material that is treated with the enzyme preparations of the invention may be manufactured of natural cellulose containing fibers or manmade cellulose containing fibers or mixtures thereof. Examples of natural cellulose are cotton, linen, hemp, jute and ramie. Examples of manmade cellulose are viscose, cellulose acetate, cellulose triacetate, rayon, cupro and lyocell. The above mentioned cellulose can also be employed as blends of synthetic fibers such as polyester, polyamide or acrylic fibers. The textile material may be yarn or knitted or woven or formed by any other means.

The cellulases of the invention, besides being especially useful for the treatment of fabric, are useful in general in any area requiring cellulase activity. In the pulp and paper industry, neutral cellulases can be used, for example, in deinking of different recycled papers and paperboards having neutral or alkaline pH, in improving the fiber quality, or increasing the drainage in paper manufacture. Other examples include the removal of printing paste thickener and excess dye after textile printing, and as a treatment for animal feed. For example, if the intended application is improvement of the strength of the mechanical pulp, then the 50K-cellulase, 20K-cellulase, 50K-cellulase B or the protein-with-CBD preparations of the invention may provide one or more of these proteins so as to enhance or facilitate the ability of cellulose fibers to bind together. In a similar manner, in the application of pulp refining, the 50K-cellulase, 20K-cellulase, 50K-cellulase B or protein-

with-CBD preparations of the invention may provide one or more of these proteins at a level that enhance or facilitate such swelling.

The invention is described in more detail in the following examples. These examples show only a few concrete applications of the invention. It is self evident for one skilled in the art to create several similar applications. Hence the examples should not be interpreted to narrow the scope of the invention only to clarify the use of the invention.

EXAMPLES

Example 1

Shake Flask and Fermentor Cultivations

For maintenance, the strains ALKO4179, ALKO4124, ALKO4237, ALKO4265 and ALKO4125 were streaked on sporulation agar (ATCC medium 5, American Type Culture Collection, Catalogue of Filamentous Fungi, 18th edition, eds., S. C. Jong and M. J. Edwards, (1991): 1 liter contains 1 g yeast extract, 1 g beef extract, 2 g tryptose, a trace amount of FeSO_4 , 10 g glucose and 15 g agar; the pH was 7.2. Agar slants were incubated at 45° for 3–6 days.

For the applications tests of ALKO4237 (Examples 3 and 4), a colony was inoculated in 500 ml of the following mineral medium (Moloney, A. P. et al., *Biotechnol. Bioeng.* 25:1169 (1983)): 1 liter contains 15 g KH_2PO_4 , 15 g $(\text{NH}_4)_2\text{SO}_4$, 2.4 ml of 1 M $\text{MgSO}_4 \times 7\text{H}_2\text{O}$, 5.4 ml 1 M CaCl_2 , 20 g Solka floc, 15 g corn steep powder, 1 g yeast extract and 10 ml 100x trace element solution 1, where 1 liter of 100x trace element solution 1 contains 0.5 g $\text{FeSO}_4 \times 7\text{H}_2\text{O}$, 0.156 g $\text{MnSO}_4 \times \text{H}_2\text{O}$, 0.14 g $\text{ZnSO}_4 \times 7\text{H}_2\text{O}$ and 0.49 g $\text{CoSO}_4 \times 7\text{H}_2\text{O}$; the pH was adjusted to pH 6.5. Cultivation was performed at 45° C. for 3 days in a rotatory shaker (250 rpm). Endoglucanase activity of about 20–25 nkat/ml was obtained.

Cellulase activity was routinely measured as endoglucanase activity according to Bailey, M. J. et al., *Enzyme Microb. Technol.* 3:153 (1981)), using 1% (w/v) hydroxyethylcellulose, HEC (Fluka AG #54290) as a substrate. The assay conditions were, if not otherwise stated, pH 7.0 and 50° C. with a 10 minute reaction time. One endoglucanase unit (1 nkat=1 ECU) is defined as the amount of enzyme that produces reducing carbohydrates having a reducing power corresponding to one nanomole of glucose in one second from HEC under the assay conditions. However, with the purified enzymes described in Examples 9–12, the assay conditions of Bailey et al., *Enzyme Microb. Technol.* 3:153 (1981) exceed the linear range, and the assay was therefore modified as described in Example 10. With every strain, the filter paper activity assay (which measures the total hydrolysis of cellulose and indicates the presence of cellobiohydrolase activity) was either under the reliable detection limit or very low.

For the determination of pH and temperature dependency (Example 2), as well as for the application tests of the strains ALKO4179, ALKO4124, ALKO4265 and ALKO4125 (Examples 3 and 4), colonies were inoculated in 500 ml of the modified thermomedium B (G. Szakacs, Technical University of Budapest, Hungary): 1 liter contained 6 g Solka floc, 6 g distiller's spent wheat grain, 3 g oat spelt xylan, 2 g CaCO_3 , 1.5 g soybean meal, 1.5 g $(\text{NH}_4)_2\text{HPO}_4$, 1 g barley bran, 0.5 g KH_2PO_4 , 0.5 g $\text{MgSO}_4 \times 7\text{H}_2\text{O}$, 0.5 g NaCl , 0.5 ml trace element solution 1 (1 liter contains: 1.6 g MnSO_4 , 3.45 g $\text{ZnSO}_4 \times 7\text{H}_2\text{O}$, and 2.0 g $\text{CoCl}_4 \times 6\text{H}_2\text{O}$) and 0.5 ml trace element solution 2 (1 liter contains: 5.0 g $\text{FeSO}_4 \times 7\text{H}_2\text{O}$ and two drops of concentrated H_2SO_4); the pH was adjusted to

pH 6.5. Cultivations were performed at 45° C. for 3 days in a rotatory shaker (250 rpm). Because in thermomedium B the endoglucanase activities of the strains ALKO4179, ALKO4124, and ALKO4237 were about 5 nkat/ml, culture filtrates were concentrated about 10 fold in an Amicon concentrator using a cut-off of 30 kDa. Endoglucanase activity obtained with ALKO 4265 was about 20 nkat/ml and with ALKO 4125 30–40 nkat/ml.

The 1 liter fermentor cultivation of ALKO4179 was performed in the following medium: 1 liter contained 10 g Solka floc, 3 g cellobiose, 4 g corn steep powder, 1.5 g $(\text{NH}_4)_2\text{HPO}_4$, 0.3 g $\text{MgSO}_4 \times 7\text{H}_2\text{O}$, 0.5 g NaCl , 2 g CaCO_3 , 0.5 ml trace element solution 1 and 0.5 ml trace element solution 2, 0.5 g KNO_3 , 0.3 g CaCl_2 , 1 g Tween 80; the pH was adjusted to pH 6.5.

The 1 liter fermentor cultivation of ALKO4124 was performed in the modified thermomedium B: 1 liter contained: 10 g Solka floc, 1 g Roth's xylan, 40 g whey, 30 g soybean meal, 2 g CaCO_3 , 5 g $(\text{NH}_4)_2\text{SO}_4$, 0.5 g KH_2PO_4 , 10 g $\text{MgSO}_4 \times 7\text{H}_2\text{O}$, 1.0 g NaCl , 1 g antifoam, 0.5 ml trace element solution 1 and 0.5 ml trace element solution 2.

The 1 liter fermentor cultivation of ALKO4237 was performed in the mineral medium mentioned above. 10% (v/v) inoculum was used. pH was maintained at $\text{pH } 6.5 \pm 0.4$ by the addition of ammonia [12.5% (v/v)] and phosphoric acid [17% (v/v)]. The fermentation temperature was 45° C. The fermentor (Biostat M, B. Braun, Melsungen, Germany) was stirred at 400 rpm and the air flow as 1 vvm. The endoglucanase activities obtained were the following: ALKO4179 about 40 nkat/ml, ALKO4124 about 90 nkat/ml and ALKO4237 about 30 nkat/ml. ALKO4265 and ALKO4125 were not cultivated in a fermentor.

ALKO4179, ALKO4124, ALKO4237 and ALKO4125 were cultivated in a 100 liter pilot fermentor in media and conditions described above. Endoglucanase activities obtained were about 40 nkat/ml with ALKO4179 and ALKO4237, about 90 nkat/ml with ALKO4124 and about 100 nkat/ml with ALKO4125. Culture filtrates were concentrated 10–20 fold in a Millipore PUF100 ultra filter and a Pellicon Us cassette concentrator using a cut-off of 10 kDa.

Example 2

Determination of the pH and the Temperature Dependence of the Endoglucanase Activities in the Culture Filtrates

For the determination of pH and temperature dependence, the strains ALKO4179, ALKO4124, ALKO4237, ALKO4265 and ALKO4125 were grown in the modified thermomedium B. Samples from the shake flask cultivations (culture filtrates) were diluted in 50 mM McIlvain's buffers (50 mM citric acid-100 mM Na_2HPO_4) of pH range 4.5–8.5. The final pH values of the culture filtrate buffer mixtures were 4.3, 5.4, 6.3, 7.3, 8.1 and 8.7 for the strain ALKO4179; 4.3, 5.4, 6.4, 7.3, 8.1 and 8.5 for the strain ALKO4124; 4.4, 5.3, 6.2, 7.1, 8.0 and 8.5 for the strain ALKO4237; 4.3, 5.4, 6.3, 7.2, 8.1 and 8.5 for the strain ALKO4265 and 4.3, 5.4, 6.4, 7.3, 8.1 and 8.5 for the strain ALKO4125. BSA was added as a protein carrier to the concentration of 100 $\mu\text{g/ml}$. Pepstatin A and phenyl methyl sulphonyl fluoride (PMSF) were added as protease inhibitors at 10 $\mu\text{g/ml}$ and 174 $\mu\text{g/ml}$, respectively. Endoglucanase activity was measured at each pH at 50° C. with 60 minutes reaction time. The endoglucanase activity of ALKO4179 exhibited more than 90% of its maximum in the pH range of about 4.5–7.5, the maximum activity was detected at about pH 5.4–6.3 (FIG. 1A).

The endoglucanase activity of ALKO4124 exhibited more than 80% of its maximum activity in the pH range about 5.5–7.5, the maximum activity was detected at about pH 6.4 (FIG. 2A). The endoglucanase activity of ALKO4265 exhibited more than 80% of its maximum activity in the pH range about 4.5–7.0, the maximum activity was detected at about pH 5.5–6.5 (FIG. 4A). The endoglucanase activity of ALKO4237 exhibited more than 80% of its maximum in the pH range of about 4.5–6.0, the maximum activity was detected at about pH 5.3 (FIG. 3A). The endoglucanase activity of ALKO4125 exhibited about 90% of its maximum in the pH range of about 4.5–7.5, the maximum activity was detected at about pH 6.5 (FIG. 5A).

For the temperature dependency determination of the endoglucanase activity, samples from the culture filtrates were diluted in 50 mM McIlvain's buffer, pH 7.0. BSA was added as a protein carrier to the concentration of 100 µg/ml. Pepstatin A and phenyl methyl sulphonyl fluoride (PMSF) were added as protease inhibitors to 10 µg/ml and 174 µg/ml, respectively. The final pH values of the culture filtrate buffer mixtures were 7.3 (ALKO4179, ALKO4124 and ALKO4125) and 7.2 (ALKO4237 and ALKO4265). Samples were incubated at 40° C., 50° C. and 60° C. for 60 minutes. The maximum endoglucanase activity of ALKO4179 was detected at 50° C. and 60° C., about 30% of the activity was retained at 40° C. (FIG. 1B). The maximum endoglucanase activity of ALKO4124 was detected at 60° C., about 70% of the activity was retained at 50° C. and 30% at 40° C. (FIG. 2B). The maximum endoglucanase activity of ALKO4237 was detected at 60° C., about 60% of the activity was retained at 50° C. and 40% at 40° C. (FIG. 3B). The maximum endoglucanase activity of ALKO4265 was detected at 60° C., about 50% of the activity was retained at 50° C. and 30% at 40° C. (FIG. 4B). The maximum endoglucanase activity of ALKO4125 was detected at 60° C., about 80% of the activity was retained at 50° C. and 70% at 40° C. (FIG. 5B).

Example 3

Indigo Dye Release in Neutral Conditions

Cellulase preparations derived from the strains ALKO4179, ALKO4124, ALKO4237, ALKO4265 and ALKO4125 (Examples 1 and 2) were tested for their ability to release dye in neutral conditions from the indigo dyed cotton-containing denim fabric to give a stone-washed look. Commercial acid cellulase product Ecoston L (Primalco Ltd, Biotec, Finland) was used as a control.

Denim fabric was obtained from Lauffenmuehl (Germany). Test fabric was prewashed 10 min at 60° C. with Ecoston A 200 (1 ml/liter, Primalco Ltd, Biotec, Finland). The fabric was then cut into 12x12 cm swatches. The colour from both sides of the fabric swatches was measured as reflectance values with the Minolta (Osaka, Japan) Chroma Meter CM 1000R L*a*b* system.

Cellulase treatments were performed in LP-2 Launder-Ometer (Atlas, Ill., USA) as follows. About 7 g of denim swatches were loaded into the 1.2 liter container that contained 200 ml of 0.05 M citrate/phosphate buffer at pH 7, or, 0.05 M citrate buffer at pH 5.2. 0.06 ml of 10% Berol 08 (Berol Nobel AS, Sweden) was added as a surfactant.

A quantity of steel balls were added into each container to help the fiber removal. Finally the cellulase solutions were added to the container as endoglucanase activity units (Example 1). The containers were then closed and loaded into a 50° C. Launder-Ometer bath. The Launder-Ometer was run at 42 rpm for 2 hours.

After removing swatches from the containers they were soaked for 10 min in 200 ml of 0.01 NaOH and rinsed for 10 min with cold water. Swatches were then dried for 1 hour at 105° C. and air dried overnight. The color from both sides of the swatches was measured with the Minolta Chroma Meter. Results from the color measurements of treated denim fabrics are shown in Table I.

TABLE I

Color Measurement of Denim Fabrics Treated with Different Cellulase Preparations.							
Source of Enzyme	ECU/g of fabric	Right side of the Fabric			Reverse side of the Fabric		
		L	b	delta E	L	b	delta E
<u>pH 7*</u>							
—	—	2.3	0.8	3.1	1.5	0.1	0.9
ALKO4237	200	6.4	3.3	7.6	2.4	1.7	3.2
	400	7.7	3.8	8.1	2.5	1.8	3.0
ALKO4179	200	5.5	2.4	6.4	2.8	1.9	3.0
	400	4.6	2.8	5.1	2.2	1.5	3.0
ALKO4124	200	4.8	2.8	6.1	3.3	1.2	2.5
	400	ND	ND	ND	ND	ND	ND
ALKO4125	200	4.0	2.7	5.6	2.3	1.5	2.3
	400	ND	ND	ND	ND	ND	ND
ALKO4265	200	2.2	3.6	5.1	-4.9	6.6	9.2
	400	ND	ND	ND	ND	ND	ND
Ecoston L	200	1.6	0.7	1.6	0	1.7	1.6
	400	1.6	0.9	1.8	-1.9	2.2	2.8
<u>pH 5.2**</u>							
Ecoston L	200	2.01	2.33	3.30	-2.74	4.35	4.71
	400	3.19	2.76	4.35	-2.56	4.83	6.71

L: Lightness unit of the fabric after the treatment minus lightness unit of the fabric before the treatment.

b: Blueness unit of the fabric after the treatment minus blueness unit of the fabric before the treatment.

delta E: Color difference in the L*a*b* color space between the specimen color and the target color (target fabric = untreated denim fabric).

ND = not done.

*the ECU activity was measured at pH 7.0.

**the ECU activity was measured at pH 4.8.

To compare the final look of the denim fabrics after washing with different cellulase preparations, the color from both sides (reverse side and right side) of the fabrics was measured. From the results shown in Table I, it can be seen that the lightness and blueness units are clearly increased on the right side of the garments washed with preparations of ALKO4179, ALKO4124, ALKO4237 and ALKO4125 cellulases, showing a good stone-washed effect. The blueness unit was also increased on the right side of the fabric washed with the ALKO4265 preparation but there was no increase in the lightness unit. This is probably because the enzyme does work at this pH but at the same time causes a lot of backstaining. There was no stone washing effect on the fabric with commercial acid product Ecoston L at pH 7 at this ECU activity.

In this study, backstaining on the reverse side of the fabric is used as an indication of the degree of backstaining on the right side of the fabric. To quantify the level of backstaining, the color was measured on the reverse side of the fabric before and after the cellulase treatment. As shown in Table I, when the ECU amounts are the same, there was practically no backstaining in the fabrics treated with the ALKO4179, ALKO4124, ALKO4237 and ALKO4125 preparations when compared to the fabrics treated with ALKO4265 or Ecoston L (pH 5.2 and 7) preparations.

Example 4

Dye Release in Neutral Conditions, No Berol

The experimental set-up was as described in Example 3 except that no Berol was used. Results from the color

measurements of treated denim fabrics are shown in Table II.

TABLE II

Color Measurement of Denim Fabrics Treated with Different Cellulase Preparations - no Berol.							
Source of Enzyme	ECU/g of fabric	Right side of the Fabric			Reverse side of the Fabric		
		L	b	delta E	L	b	delta E
pH 7*							
—	—	2.1	0.5	2.2	1.7	-1.1	2.0
ALKO4237	200	5.5	3.1	7.0	1.8	2.3	3.5
ALKO4179	200	4.4	3.2	5.6	1.4	2.2	2.7
ALKO4124	200	4.2	2.9	5.0	1.1	2.0	2.4
ALKO4125	200	3.5	2.6	4.4	1.6	1.4	2.5
ALKO4265	200	3.3	3.3	5.3	-5.7	6.6	10.0
Ecoston L	200	1.4	0.9	1.7	0.3	1.4	1.8
	400	1.4	0.8	1.7	-0.1	1.7	1.8
pH 5.2**							
Ecoston L	200	2.0	2.1	2.9	-4.0	4.8	5.4

L: Lightness unit of the fabric after the treatment minus lightness unit of the fabric before the treatment.

b: Blueness unit of the fabric after the treatment minus blueness unit of the fabric before the treatment.

delta E: Color difference in the L*a*b* color space between the specimen color and the target color (target fabric = untreated denim fabric).

ND = not done.

*the ECU activity was measured at pH 7.0.

**the ECU activity was measured at pH 4.8.

When compared with results obtained with the inclusion of Berol (Example 3), the data in Table II show that almost the same stone-washing effect can be achieved with the ALKO4179, ALKO4124, ALKO4237 and ALKO4125 cellulase preparations in the absence of the helping agent Berol.

Example 5

Backstaining in Denim Wash with Different Cellulases

In the literature, it is reported that backstaining is dependent on pH and/or the type of enzyme. However, as shown herein, it was found that backstaining depends only indirectly on pH (FIGS. 6A and 6B and 7A and 7B).

Two neutral cellulase preparations from ALKO4237 and from ALKO4265 and acid cellulase product Ecoston L were studied in small scale denim wash with an equal enzyme dosage at pH 5 and pH 7. The stonewash effect was determined by measuring the increase of lightness and blueness as reflectance units on the right side of the fabric and backstaining (redeposition of indigo on the surface of

fibers) was determined as blueness increase and lightness decrease on the reverse side. At pH 7, the neutral cellulases from ALKO4237 caused a clear increase in lightness and blueness on the right side and no backstaining was observed (FIGS. 6A and 6B). A similar stonewash effect was found at pH 5 but with a slight backstaining. At pH 7, the other neutral cellulase, ALKO4265, brightened blueness on the right side but backstained intensively on the reverse side. At pH 5 similar effects were obtained with both ALKO4265 and ALKO4237 preparations. At pH 7, the acid cellulase did not backstain or impart a lightness on the right side (when using similar endoglucanase activity dosages as with ALKO4265 and ALKO4237, FIGS. 7A and 7B, 1xdosage), probably because it did not work at this pH. On the other hand, at pH 5, lightness and blueness were increased on the right side and backstaining was clearly perceptible on the reverse side. Based on these results, backstaining can occur at both pH values depending on the cellulase preparation used.

Example 6

Use of the Neutral Cellulase-Containing Enzyme Preparations in Biofinishing of Cotton-Containing Woven Fabric

100% cotton woven fabric was subjected to treatment with ALKO4237 (Example 1) and ALKO4467 cellulases in Launder-Ometer. ALKO4467 is a UV-mutant with higher cellulase activity derived from ALKO4125.

100% cotton woven fabric (obtained from Pirkanmaan Uusi Värjäamo Ltd) was pretreated as in Example 7. The cellulase treatment conditions were as described in Example 3 except that no Berol was used and the liquid ratio was 1:15 (volume of liquid per weight of fabric). Cellulases were dosed as ECU activity units (Example 1).

The following methods were used for evaluation of the effect of the enzyme preparations in biofinishing of cotton fabric: Weight loss of the treated fabrics was defined as percentage from weight of the fabric before and after the test (before weighing the fabrics were conditioning in a atmosphere of 21±2° C. and 50±5% RH). Evaluation of the surface cleaning effect of the enzyme treated fabrics was performed by a panel consisting of three persons. The fabrics were ranked on a score from 1 to 5, where 5 gave a clean surface. The Martindale Rubbing method (SFS-4328) was used for evaluation of pilling. Pilling was evaluated by a panel after 200 and 2000 cycles of abrasion (1=many pills, 5=no pills).

In Table III is shown that treatment of the cotton fabric with ALKO4237 and ALKO4467 cellulase preparations results in a good surface cleaning and marked reduction in the pilling tendency at both pH 5 and 7.

TABLE III

Weight loss, surface cleaning effect and pitting tendency of the cotton fabrics treated with neutral cellulases in Launder-Ometer.

preparation	dosage ECU/g	time h	pH	weight loss %	surface cleaning effect	pilling 200 cycles	2000 cycles
—	—	1	5	0	1.0	1.0	1.0
ALKO4237	200	1	5	2.3	3.5	4.0	3.8
ALKO4237	400	1	5	3.2	3.5	4.0	3.8
ALKO4467	200	1	5	1.2	2.5	3.7	3.4
ALKO4467	400	1	5	1.9	2.8	3.7	3.4
—	—	2	5	0.1	1.0	1.0	1.0
ALKO4237	200	2	5	4.4	4.0	4.2	4.1
ALKO4237	400	2	5	6.0	4.3	4.2	4.3

TABLE III-continued

Weight loss, surface cleaning effect and pilling tendency of the cotton fabrics treated with neutral cellulases in Launder-Ometer.							
preparation	dosage ECU/g	time h	pH	weight loss %	surface cleaning effect	pilling 200 cycles	2000 cycles
ALKO4467	200	2	5	3.0	3.5	4.0	3.8
ALKO4467	400	2	5	4.0	3.8	4.0	3.9
—	—	1	7	0	1.0	1.0	1.0
ALKO4237	200	1	7	2.5	3.0	3.7	3.5
ALKO4237	400	1	7	3.8	4.0	4.0	3.9
ALKO4467	200	1	7	0.8	2.0	3.5	3.3
ALKO4467	400	1	7	1.4	2.0	3.6	3.7
—	—	2	7	0.1	1.0	1.2	1.1
ALKO4237	200	2	7	4.8	4.0	3.8	4.0
ALKO4237	400	2	7	6.0	4.3	4.0	4.3
ALKO4467	200	2	7	2.2	2.5	4.0	3.4
ALKO4467	400	2	7	3.0	3.3	3.8	3.7

Example 7

Use of the Neutral Cellulase-Containing Enzyme Preparations of the Invention in Biofinishing

7a. Use of Enzyme Preparations in the Biofinishing of Woven Fabric and Knit

100% cotton woven fabric or 100% cotton knit are subjected to treatment with the cellulases of the invention (Example 1) in a semi-industrial drum washer (Esteri 20 HS-P). The treatment conditions are as follows:

A. Pretreatment (only for woven fabrics)
60° C., 10 minutes, Ecoston A200 (Primalco Ltd, Biotec, Finland) 1 ml/l water.

B. Enzyme treatment
temperature 50–60° C., pH 7;
liquid ratio 5–20:1 (volume of liquid per weight of fabric);
treatment time 20–90 minutes, preferably 30–60 minutes;
and

enzyme dosage 50–900 nkat/g fabric or knit, preferably 200–600 nkat/g fabric or knit.

C. "After-washing" treatment
40° C., 10 minutes, alkaline detergent

D. Drying treatment

The following standard methods are used for evaluation of the surface cleaning effect of enzyme preparations: The Martindale Rubbing Method (SFS-4328) and the Laundering Durability Test (SFS-3378). Treatment with the cellulase preparations of the invention results in a surface cleaning effect, an improvement in the softness and smoothness of the fabric and knit and a reduction in the pilling tendency.

7b. Use of Enzyme Preparations in the Finishing of Lyocell Fabrics and Knits

The cellulase preparations of the invention can be used in fibrillation control and different finishing processes of 100% lyocell fabrics and knits and blends thereof. The following treatment conditions in semi-industrial drum washer (Esteri 20 HS-P) are used in order to create the peach effect on lyocell fabric:

A. Sodium carbonate 2.5 g/l; 60° C., treatment time of 60 minutes;

B. Rinse;

C. Enzyme treatment: temperature of 50–60° C., pH 7, liquid ratio 5–20:1, treatment time 40–120 minutes, preferably 45–90 minutes, and an enzyme dosage of 100–1500 nkat/g fabric, preferably 400–800 nkat/g fabric;

20

D. Aftertreatment: Alkaline detergent wash at 40° C. for 10 minutes;

E. Rinse; and

F. Dry.

25

The result is a peach skin effect.

Example 8

Use of Enzyme Preparations in Biostoning

Denim garments were subjected to treatment with the neutral cellulase preparations (Example 1) in a semi-industrial drum washer (Esteri 20 HS-P) to give the garments a stonewashed appearance. About 1.0 kg of denim garments (contained two different kinds of fabric) were used per machine load.

30

The treatment conditions were as follows.

A. Desizing. 100 liters water, 60° C., 10 minutes; 100 ml Ecoston A200 (Primalco Ltd, Biotec, Finland).

B. Cellulase Treatment 100 liter water, 50° C., 45 minutes; 10 g Berol 08 (Berol Nobel AS, Sweden); 30 g citric acid+128 g Na₂HPO₄×2 H₂O to give pH 7.

35

Neutral cellulase preparations were dosed as endoglucanase activity units (ECU, Example 1):

1. ALKO4237, 260 ECU/g of garment

2. ALKO4179, 260 ECU/g of garment

3. ALKO4124, 300 ECU/g of garment

4. ALKO4125, 250 ECU/g of garment

40

C. Afterwashing. Alkaline detergent wash, 40° C., 10 minutes.

D. Drying.

45

The results were evaluated by visual appearance of the garments and by measuring the color as reflectance values with the Minolta Chroma Meter CM 1000R L*a*b system (Table IV). A good stonewashed effect was obtained with all these cellulase-treated garments. No backstaining (examined on the inside of the garment) could be seen visually in any of these cellulase-treated garments.

50

From the results of the color measurements shown in Table IV, it can be seen that the lightness and blueness units are clearly increased on the outside of the garments washed with the neutral cellulase preparations, showing a good stonewashed effect.

55

60

65

TABLE IV

Color Measurement of Denim Garments with Different Cellulase Preparations				
Source of	Outside of the garment		Inside of the garment	
Enzyme	L	b	L	b
A. Fabric 1				
untreated	24.1	-8.5	57.1	0.17
washed without cellulase	21.4	-14.0	54.5	-4.3
ALKO4237	26.7	-17.3	56.5	-4.9
ALKO4179	26.8	-17.0	56.3	-4.5
ALKO4125	28.0	-17.4	57.8	-4.1
ALKO4124	26.4	-17.5	57.1	-4.8
B. Fabric 2				
untreated	22.5	-8.3	57.6	0.66
ALKO4237	25.0	-16.3	56.1	-4.3
ALKO4179	25.0	-15.8	55.4	-4.4
ALKO4125	26.7	-17.0	56.8	-4.0
ALKO4124	25.6	-17.0	56.4	-4.0

L = Lightness unit of garment after the treatment (the higher the value, the lighter the garment).
b = Blueness unit of garment after the treatment (the more negative value, the more bluing in the garment).

or than the unfractionated concentrate. A mixture of Peaks I and II containing 70 ECU of each/g denim was also tested. This resulted in an L (right) value of 7.3 and b (reverse) of 2.5. Thus, this mixture was more effective than either peak alone.

The purification procedure was scaled up to obtain homogenous samples of some of the desired proteins in these peaks. Concentrated ALKO4237 growth medium (4.5 liters) was fractionated with ammonium sulphate. The proteins that precipitated between 17 g and 42 g of ammonium sulphate per 100 ml of concentrate were suspended in 0.9 liter of 25 mM Tris/HCl pH 7.2 containing 0.25 mM EDTA and then diluted with water to a conductivity of 4 mS/cm and adjusted with 1M NaOH to pH 8.0. The resulting solution (about 45 liters) was pumped at 150 ml/min through a 6.3 liter column of DEAE-Sepharose FF™ at room temperature. Peak I endoglucanase activity did not bind under these conditions. Bound proteins were eluted at 110 ml/min with a linear gradient from 0.0 to 0.5 M NaCl in 20 liters of 25 mM Tris/HCl pH 7.7 containing 0.25 mM EDTA. Peak II endoglucanase eluted at about 14 mS/cm. Instead of the separate Peaks III and IV seen with small scale separations in DEAE in the cold room, a single peak, called Peak III/IV, eluted at about 25 mS/cm.

TABLE V

Indigo Dye Release by DEAE-Sepharose™ Pools in Neutral Conditions

	ADDITION									
	None	Concentrate	Peak I	Peak II	Peak III	Peak IV				
ECU/g	0	100	200	310	185	340	97	260	95	190
mg/g	0	10	20	41	9	26	24	46	5	10
L(right)	2.9	5.2	7.0	5.5	7.3	10.3	4.4	5.8	3.9	4.3
b(reverse)	0.4	2.6	3.5	3.5	2.9	2.5	0.9	1.4	0.1	0.5

The parameter L (right) indicates the lightening of the right side of the blue denim, and b (reverse) indicates the bluing of the reverse side (i.e., backstaining). The fabric was washed in the LP-2 Launder-Ometer and then measured with the Minolta ChromaMeter, as described in Example 3, except that no Berol was used and the buffer that was used was 0.05M McIlvaine pH 7 (see Data for Biochemical Research, Dawson, R., et al., eds., 1969, Oxford Univ. Press). The dosage is shown as both ECU/g of denim and mg protein/g of denim.

Example 9

Purification of Neutral Cellulases

Concentrated growth medium from ALKO4237 was fractionated at 7° C. on DEAE Sepharose CL6B with a linear gradient from zero to 0.5 M NaCl in 25 mM Tris/HCl pH 7.2. Four peaks of endoglucanase activity at pH 4.8 were found. Peak I, containing about 10% of the recovered ECU, eluted at about 150 mM NaCl, Peak II (about 30% of ECU) at 230 mM NaCl, Peak III (about 20% of ECU) at 270 mM NaCl and Peak IV (about 40% of ECU) at 320 mM NaCl. Table V shows the results when these peaks were tested for their utility in biostoning at neutral pH and 50° C.

These results show that on both an ECU basis and a total protein basis, Peak II was more effective than any other peak

45

Proteins in Peak II (3.5 liters) were precipitated with ammonium sulphate (450 g/liter) and suspended in 170 ml 25 mM PIPES/KOH pH 6.0 containing 1 mM EDTA. Portions of this material were transferred to 25 mM sodium acetate pH 4.0 containing 1 mM EDTA by gel-filtration on a 5x29 cm column of G25 Sephadex™ (coarse) and then fractionated on SP-Sepharose™. FIG. 8 shows the result that was obtained when 11.7 g of these proteins was applied to a 4.5x31 cm column of SP-Sepharose™ in 25 mM sodium acetate pH 4.0 containing 1 mM EDTA at 150 ml/h and the column developed at 75 ml/h with a linear gradient from 0.0 to 0.4 M NaCl in 3.4 liters of the same buffer. Most of the endoglucanase eluted at 0.2 M NaCl. The modified assay described in Example 10 was used. When active fractions were stored at 7° C., a crystalline precipitate appeared in them and contained nearly all the endoglucanase activity. Active fractions (15 ml) in which crystallization was slow, were induced to form crystals by seeding with 30 µl of suspension from fractions already containing crystals. After 2 to 3 days, the crystals were collected by centrifugation, washed with 25 mM PIPES/KOH pH 6.0 containing 1 mM EDTA and dissolved in 25 mM Tris/HCl pH 7.2 containing

65

0.25 mM EDTA. Analysis by SDS-PAGE showed the washed crystals contained a virtually homogenous protein with an apparent molecular mass close to 20 kDa (the error in SDS-PAGE estimations of molecular mass is at least $\pm 10\%$, and may be much greater for unusual proteins). This protein is called the 20K-cellulase. Contaminating protein could also be removed by gel-filtration on G50 Sephadex™ in 50 mM PIPES/KOH pH 6.0 containing 1 mM EDTA. An example of this is shown in FIG. 9, where unwashed crystals were purified by gel-filtration. The endoglucanase activity co-eluted with the 20 kDa protein well after the cytochrome c (11.2 kDa) volume, showing that this 20 kDa protein is abnormally retarded by interaction with Sephadex™.

Proteins in Peak III/IV were precipitated with ammonium sulphate and transferred to 25 mM sodium acetate pH 4.0 containing 1 mM EDTA in the same way as described for the Peak II proteins. Upon transfer to 25 mM sodium acetate pH 4.0, a large precipitate formed and was discarded. The active supernatant was fractionated on SP-Sepharose™. At low protein loading (e.g. 200 mg protein to a 2.5x11 cm column as shown in FIG. 10, most of the endoglucanase activity bound to the column and was eluted with a NaCl gradient at about 50 mM NaCl. This active peak was followed by a second peak of inactive protein.

SDS-PAGE analysis showed that the active and inactive peaks both contained several proteins, including proteins with apparent molecular masses close to 50 kDa that could not be distinguished from each other by SDS-PAGE. Both peaks were further purified by chromatography on Phenyl Sepharose™.

The active fractions (fractions 15 to 18 in FIG. 10) were pooled, adjusted to 50 mM PIPES/KOH pH 6.0 (by addition of 0.25 M PIPES/KOH pH 6.0) and 15 g % ammonium sulphate (by addition of solid ammonium sulphate) and applied to a 1.5x8.5 cm column of Phenyl Sepharose™ equilibrated with 25 mM PIPES/KOH pH 6.0 containing 1 mM EDTA and 15 g % of ammonium sulphate. The column was developed with a linear gradient from 15 to 0 g % ammonium sulphate in 104 ml of 25 mM PIPES/KOH pH 6.0. After the end of the gradient, the column was further washed with 25 mM PIPES/KOH pH 6.0. Two protein peaks eluted on the gradient, first a small peak of inactive protein and then a major peak containing most of the endoglucanase activity. SDS-PAGE analysis (FIGS. 11A and B) showed that both peaks contained essentially homogenous proteins with apparent molecular masses close to 50 kDa (i.e., they migrate slightly slower than the BioRad prestained ovalbumin standard, which has an apparent molecular mass of 47 kDa). These two proteins could not be distinguished by the inventors' SDS-PAGE analyses, even when they were run together as mixtures. The protein in the active peak was called 50K-cellulase and the protein in the inactive peak was called 50K-protein B. Larger amounts of 50K-cellulase B were obtained by fractionation of the second (and inactive) peak eluted from SP-Sepharose™ (fractions 19 to 23 in FIG. 10) on Phenyl Sepharose™ in exactly the same way as described above for the active fractions.

Production of still larger amounts of 50K-cellulase and 50K-cellulase B was facilitated by overloading the SP-Sepharose™ column. For example, when 15 g of protein was applied to a 4.5x31 cm column of SP-Sepharose™, instead of binding to the column, the 50K-cellulase was apparently displaced by more strongly bound proteins, and eluted before the NaCl gradient. This material was already highly purified, and homogenous 50K-cellulase was isolated from it by chromatography on Phenyl Sepharose™ as described above.

In order to speed up the purification of larger amounts of 50K-cellulase the SP-Sepharose and Phenyl Sepharose columns were reversed. After adjusting the ammonium sulphate concentration to about 15 g %, the proteins precipitated in Peak III/IV were applied into Phenyl Sepharose as described before. With high overloading (e.g. 17 g of protein applied to a 3.2x25 cm column of Phenyl Sepharose) most of the total protein ran through the column, but 50K-cellulase (containing most of the endoglucanase activity) was bound and eluted at the end of linear gradient from 15 to 0 g % of ammonium sulphate in 25 mM PIPES/KOH pH 6.0. Western analysis with a rabbit antiserum recognizing 50K-cellulase B showed that the 50K-cellulase B eluted just before 50K-cellulase. Further purification was achieved by fractionation on SP-Sepharose as described earlier. In this reversed order of SP-Sepharose and Phenyl Sepharose the proteins in Peak III/IV precipitated with ammonium sulphate could be applied directly to the next purification step without removing salt. The large protein precipitate, which appeared upon transfer of the concentrated proteins in Peak III/IV directly into 25 mM sodium acetate pH 4.0 for SP-Sepharose, could also be avoided this way. As the 50K-cellulase only just binds to SP-Sepharose, the preceding fractionation on Phenyl Sepharose markedly reduced the apparently interfering total protein load on SP-Sepharose.

50K-cellulase and 50K-cellulase B were each tested in the Launder-Ometer to see if they are responsible for the beneficial effects of Peak IV reported in Example 10. Both proteins were found to have beneficial effects (Table VI). At the low concentrations used in this experiment, they did not themselves increase the release of indigo dye from the outer face of the denim (i.e., L_{right} did not increase) but they effectively decreased the back-staining of dye onto the inner face of the denim ($L_{reverse}$ became more positive and $b_{reverse}$ became smaller) especially when used together with 20K-cellulase.

The 20K-cellulase performed well in Launder-Ometer tests at pH 5 as well as at pH 7. At pH 5, 0.2 mg of 20K-cellulase per g of denim increased L_{right} from 3.2 to 5.2. Addition of 50K-cellulase at 0.1 mg per gram of denim together with the 20K-cellulase also decreased the back-staining at pH 5 ($L_{reverse}$ and $b_{reverse}$ changed from 0.0 and 2.6 with 20K-cellulase alone to 1.3 and 1.5, respectively, with the mixture of 2° K and 50K-cellulases).

TABLE VI

Indigo Dye Release by 20K-cellulase, 50K-cellulase and 50K-cellulase B
Conditions were the same as in Table V. The dose is shown as mg protein per gram of denim.

Sample	Dose (mg/g)	L_{right}	$L_{reverse}$	$b_{reverse}$
Buffer alone	—	2.8	-0.6	1.6
20K-cellulase	0.18	5.6	-1.0	4.0
	0.09	4.8	-1.5	3.3
50K-cellulase	0.15	2.6	-0.3	1.0
	0.075	3.0	0.4	1.3
50K-cellulase B	0.31	2.8	1.3	0.8
	0.15	2.7	1.5	0.5
20K-cellulase + 50K-cellulase	0.18 + 0.075	5.6	0.3	2.5
	0.09 + 0.075	5.1	0.3	2.1
20K-cellulase + 50K-cellulase B	0.18 + 0.15	4.7	0.0	3.0

Properties of the 20K-cellulase

Although polyclonal antibodies prepared against cellulases purified from *Trichoderma reesei* (designated anti-EGI, anti-CBHI and anti-CBHII antibodies) recognized proteins in the ALKO4237 growth medium, there was only a very weak cross-reaction with pure 20K-cellulase under the same conditions of Western blot analysis.

When growth medium from ALKO4237 was probed on Western analysis with antiserum raised in rabbits against pure 20K cellulase, a strong band at about 35 kDa was observed in addition to the 20 kDa band. No apparent endoglucanase activity could be detected for this 35 kDa protein. Also, a weaker band was seen immediately ahead of the 20 kDa band (FIG. 14).

ALKO4124 gave an almost identical pattern as ALKO4237, indicating that this and other fungi probably contain cellulases very similar to the 20K-cellulase of the present invention.

Amino acid sequences of tryptic peptides derived from 20K-cellulases are shown in FIG. 17 #429 corresponds to SEQ ID NO: 1, #430 corresponds to SEQ ID NO: 2, #431 corresponds to SEQ ID NO: 3, #432 corresponds to SEQ ID NO: 4, #433 corresponds to SEQ ID NO: 5, #439 corresponds to SEQ ID NO: 6, fr 9 corresponds to SEQ ID NO: 7, fr 14 corresponds to SEQ ID NO: 8, fr 16 corresponds to SEQ ID NO: 9, fr 17 corresponds to SEQ ID NO: 10, fr 28 corresponds to SEQ ID NO: 11, and fr 30 corresponds to SEQ ID NO: 12.

Purified 20K-cellulase performed well in biostoning at neutral pH without the addition of other enzyme activities as shown in Table VII.

TABLE VII

Biostoning by Purified 20K-cellulase						
Conditions were as in the experiment shown in Table V. Dosage is shown as mg protein/g denim fabric. "Whole medium" indicates the unfractionated ALKO4237 concentrated growth medium.						
Addition	Dosage	L _{right}	b _{right}	L _{reverse}	b _{reverse}	
Buffer	0.0	3.6	0.1	0.5	0.6	
20K	0.72	8.9	2.9	-1.1	4.7	
20K	0.25	6.0	2.3	-0.5	3.6	
20K	0.07	5.3	1.7	-0.4	2.9	
Whole medium	20	6.1	2.8	-2.9	5.5	

Compared to the unfractionated medium, 20K-cellulase resulted in the same degree of lightening (L_{right} =6.0-6.1) at 1/80th the protein dosage. Further, there was less backstaining onto the reverse side face of the fabric ($L_{reverse}$ =-0.5 compared to -2.9 and $b_{reverse}$ =3.6 compared to 5.5). Fabric treated with 20K-cellulase had an agreeable soft texture.

Although 20K-cellulase performed surprisingly well without other additions, even better fabric appearance and texture resulted when 20K was used together with the DEAE-Sepharose pools I, III or IV (Table VIII).

TABLE VIII

Synergy in Biostoning Between 20K-cellulase and Endoglucanase Pools Eluted from DEAE-Sepharose						
Conditions were as in Table V.						
Addition	Dosage	L _{right}	b _{right}	L _{reverse}	b _{reverse}	
Buffer	0.0	3.8	0.2	-0.7	1.5	
20K	0.18	5.8	2.3	-2.2	5.5	
Pool I	15	5.1	1.9	-3.1	5.7	
Pool III	47	5.2	1.6	-0.1	2.6	
Pool IV	14	5.6	0.9	0.4	1.8	
20K + Pool I	15.18	7.1	2.8	0.7	3.3	
20K + Pool III	47.18	7.6	3.1	-1.7	5.3	
20K + Pool IV	14.18	8.6	2.6	0.8	3.2	
Whole medium	20	5.7	2.4	-4.1	5.9	

The mixtures of 20K-cellulase with Pools I, III and IV caused more lightening (increased L_{right}) than either component alone. At least for the combination of 20K-cellulase with Pool IV, it is clear that this is because of synergy and not merely an additive effect. Further, the backstaining with all mixtures was actually less ($L_{reverse}$ more positive, $b_{reverse}$ less) than the backstaining observed with 20K-cellulase alone. The combination of 20K with Pool IV was particularly effective. Pool IV contains many proteins, one of which (a 50 kDa polypeptide) copurifies with endoglucanase activity during chromatography of Pool IV on Sephadex G100 and S-Sepharose. While good biostoning is achieved with 20K-cellulase alone, better results are possible with 20K-cellulase plus one or more proteins purified from Pool IV. Biostoning with mixtures of the 20K-cellulase and the 50K-cellulase and the 50K-cellulase B purified from Pool III/IV have already been presented (Table VI in Example 9). Therefore, the present invention is not limited to the use of only the 20K-cellulase. Other proteins in the ALKO4237 medium are useful alone or in suitable combinations.

In the standard endoglucanase assay described by Bailey et al. (1981, loc. cit.), the enzyme amount is chosen that produces, in 10 min and pH 4.8 (0.05 M Na-citrate buffer), about 0.6 mM reducing equivalents from 1% hydroxyethylcellulose, resulting in a final absorbance change (ΔA_{540}) of between 0.2 and 0.25. This far exceeds the range in which ΔA_{540} is proportional to the amount of 20K-cellulase.

Therefore, the procedure was modified as follows. Enough enzyme was used to produce about 0.2 mM reducing equivalents in 10 min in 0.05 M HEPES buffer (pH 7.0). To reach the threshold concentration of reducing equivalents above which color is formed in the DNS system, 0.12 mM glucose was freshly added to the stock DNS reagent. This method (called the "modified" method) was used when characterizing the endoglucanase activity of the 20K-cellulase and also the 50K-cellulase. With 1% hydroxyethylcellulose as substrate, the range in which ΔA_{540} is proportional to the amount of 20- and 50K-cellulase is relatively narrow, and so 2% carboxymethylcellulose was taken as an alternative substrate. With 2% carboxymethylcellulose, the range of linear correlation between ΔA_{540} and the amount of 20K- and 50K-cellulase was broader than with 1% hydroxyethylcellulose. The endoglucanase activity determined with 2% carboxymethylcellulose was about 8-10-fold for 20K-cellulase and about 50-fold for 50K-cellulase compared with that determined with 1% hydroxyethylcellulose.

No activity of 20K-cellulase was detectable for 4-methylumbelliferyl- β -D-lactoside, a characteristic sub-

strate of cellobiohydrolases. The activity towards filter paper was also very low, but detectable.

The 20K-cellulase was relatively heat stable. It was incubated at 7 μ g/ml and 100° C. in 25 mM Tris-HCl, 0.2 mM EDTA, for 30 or 60 min. and then assayed at pH 7.0 and 50° C. 52% and 35% respectively, of the endoglucanase activity remained at pH 7.2. 40% and 22%, respectively, remained at pH 8.8. (These pH values were measured at room temperature; the actual pH at 100° C. is somewhat lower.) At 80° C., pH 7.2, 70% of the activity remained for 60 min.

These results indicate that the enzyme is suitable for applications in which it may be (e.g., accidentally) exposed to elevated temperatures. As well as being resistant to irreversible inactivation at high temperatures, the enzyme exhibited an optimum temperature of 70° C. during 10 min. assays at pH 7.0 (FIG. 15). The decreased activity observed above 70° C. was mainly due to a reversible change in enzyme conformation: the enzyme recovered most of its activity when returned to 50° C.

At 50° C., the 20K-cellulase exhibited 80% or more of its maximum activity throughout the pH range 4 to 9, and nearly 50% at pH 10. This was the case in both 10 min. (FIG. 16A) and 60 min. (FIG. 16B) assays. These figures also show the pH dependence of the enzyme at 70° C. With 10 min. assays, the enzyme was more active at 70° C. than it was at 50° C. over the range pH 4.5 to 8 and about equally active at pH 10 (FIG. 16A). With 60 min. assays (i.e., approaching commercial conditions), the enzyme was more active at 70° C. than it was at 50° C. between pH 5.5 and 7.5. However, it was only slightly less active at 70° C. than at 50° C. up to pH 10. In practice, this means that the enzyme can be used equally well over a wide range of pH and at temperatures up to at least 70° C.

Example 11

Properties of the 50K-cellulase

Pure 50K-cellulase had both endoglucanase activity (against hydroxyethylcellulose) and cellobiohydrolase activity (against 4-methylumbelliferyl- β -D-lactoside, assayed essentially as described by van Tilbeurgh et al, in *Methods in Enzymology* [1988] vol. 160, pp 45-59). A sample of the pure enzyme with an A_{280} of 1.8 contained 2030 ECU/ml and 300 PCU/ml at pH 7.0 and 50° C. (one PCU is the amount of activity that liberates 1 nmol of methylumbelliferone per second).

In Western analyses, 50K-cellulase was strongly recognized by antiserum (KH 1057) raised against endoglucanase I of *T. reesei*, but only weakly by antisera (KH 1050 and KH 1053, respectively) against cellobiohydrolases I and II of *T. reesei*. It was not recognized by the antiserum raised against 20K-cellulase (FIG. 14). When the growth medium of ALKO 4237 was probed in Western analyses with rabbit antiserum raised against 50K-cellulase itself, only one obvious band (which had a molecular mass between 33 and 47 kDa) was seen in addition to the very strong band at about 50 kDa.

The apparent molecular mass of 50K-cellulase by SDS-PAGE decreased by about 2 to 5 kDa when the protein was treated with endoglycosidase H_1 , indicating that the enzyme contains carbohydrate removable by this endoglycosidase.

50K-cellulase was unusually resistant to tryptic digestion, indicating that it has an unusually stable structure. However, it was cleaved by treatment with cyanogenbromide, and the resulting fragments could then be digested with trypsin or

with lysylendopeptidase C. Sequences of some of the peptides so obtained are shown in Table IX.

TABLE IX

Sequences of peptides isolated from the 50 K-cellulase (uncertain residues in lower case)	
#507 (SEQ ID NO:13)	VYLLDETEHR
#509 (SEQ ID NO:14)	XXLNPGGAYYGT
#563 (SEQ ID NO:15)	MeEGAECEYDGVCDKDG
#565 (SEQ ID NO:16)	NPYRVXITDYYGNS
#603 (SEQ ID NO:17)	DPTGARSELNPGGAYYGTGYXDAQ
#605 (SEQ ID NO:18)	XXVPDYHQHGVda
#610 (SEQ ID NO:19)	NEMDIXEANSRA
#611 (SEQ ID NO:20)	LPXGMNSALYLSEMDPTGARSELNP
#612 (SEQ ID NO:21)	VEPSPEVTYSNLRXGEIXqXF
#619 (SEQ ID NO:22)	DGCGWNPYRVvITtDYyN
#620 (SEQ ID NO:23)	LPCGMXSALY
#621 (SEQ ID NO:24)	ADGCQPRNTYIVLDdLIHPXXQ

The 50K-cellulase is a stable enzyme that exhibits endoglucanase activity over a wide range of pH values and at high temperatures, so it is suitable for use in many industrial conditions. At pH 7.0 and with 60 min reaction times, it has an optimum temperature between 65 and 70° C., and even with this long reaction time still exhibits, at 75° C., 50% of the activity observed at 50° C. (FIG. 12).

With 60 min reaction times, the pH optimum was very broad at 50° C., with essentially constant activity between pH 4.4 and 7.0, and activities at pH 9 and 10 equal to 50% and 30%, respectively, of that at pH 7.0. At 70° C., there was a clear optimum at pH 6, and, between pH 5 and 7, the activity (with 60 min reaction times) was 3-fold or more greater than that at 50° C. However, at pH 4.4 and pH values above 8, the activity was greater at 50° C. than at 70° C. (in 60 min assays), suggesting that the stability of the enzyme decreases at 70° C. right side the pH range 5 to 7.5. The pH-dependence is illustrated in FIG. 13.

Example 12

Properties of 50K-cellulase B

No detectable endoglucanase activity could be measured for the 50K-cellulase B (previously called 50K-protein B) with hydroxyethylcellulose or carboxymethylcellulose. At acidic pH, the 50K-cellulase B had a low cellobiohydrolase activity, which (measured with 4-methylumbelliferyl- β -D-lactoside) at pH 5 was less than 0.1% of that of the 50K cellulase. In addition, the 50K-cellulase B had a detectable activity towards filter paper at pH 4.8 and acid swollen, amorphous Solca Floc-cellulose at pH 5 and 7 used in enzyme activity determinations.

In Western analyses, 50K-cellulase B was strongly recognized by antiserum (KH1050) raised against cellobiohydrolase I of *T. reesei*, but only weakly by antisera against cellobiohydrolase II or endoglucanase I of *T. reesei* or against the 50K-cellulase. It was not recognized by antiserum raised against the 20K-cellulase (FIG. 14). Table X shows sequences of peptides isolated from 50K-cellulase B.

TABLE X

Sequences of peptides isolated from the 50 K-cellulase B (uncertain residues in lower case)	
#534 (SEQ ID NO:25)	vGNPDFYgK
#535 (SEQ ID NO:26)	FGPIGSTY
#631 (SEQ ID NO:27)	LSQYFIQDGERK

TABLE X-continued

Sequences of peptides isolated from the 50 K-cellulase B (uncertain residues in lower case)		
#632	(SEQ ID NO:28)	FTVVS ^r FEENK
#636	(SEQ ID NO:29)	HEYGTNVGSRFYLMNGPDK

Example 13

Stability of Neutral Cellulases in Different Detergents

Stability of the neutral cellulase preparations were tested in three different detergent solutions. The detergent solutions were OMO® Total (or OMO® Neste, Lever UK), OMO® Color (Lever S.A.) and Colour Detergent Liquid (Unilever, The Netherlands). The tested cellulase preparations were ALKO4125, ALKO4179, ALKO4237 and ALKO4265 (Example 1) concentrated culture filtrates and purified 20K- and 50K-cellulases from the ALKO4237 strain (Example 9).

Cellulase preparations were incubated at 40° C. in 0.25% detergent solutions. The activity against hydroxyethylcellulose (ECU/ml, Example 1) was measured (pH 7, 50° C.) from samples taken after 5–30 minutes incubation.

The tested preparations were as follows:

Culture filtrates:

ALKO4125: 780 ECU/ml (pH 7, 50° C.)

ALKO4179: 830 ECU/ml

ALKO4265: 760 ECU/ml

ALKO4237: 650 ECU/ml

Purified proteins:

20K-cellulase: 9423 ECU/ml

50K-cellulase: 10100 ECU/ml

The results are shown in Tables XI–XIII.

ALKO4179, ALKO4265 and ALKO4237 cellulase preparations and 20K- and 50K-cellulases stay almost 100% stable for 30 minutes at 40° C. in all three tested detergents. ALKO4125 stays stable for 30 minutes at 40° C. in Colour Detergent Liquid and in OMO® Neste.

TABLE XI

Stability of different cellulases in 0.25% Colour Detergent Liquid (pH 7.5-7.9).							
preparation	enzyme dosage	pH*	% of activity left				
			0'	5'	10'	20'	30'
<u>Culture filtrates:</u>							
ALKO4125	6	7.3	100	97	98	98	99
ALKO4179	6	7.1	100	99	100	100	10
ALKO4265	6	7.2	100	100	100	100	100
ALKO4237	6	7.1	100	100	82	95	100
<u>Purified proteins from ALKO4237:</u>							
20K-cellulase	1	7.8	100	98	99	97	100
50K-cellulase	1	7.6	100	100	100	100	100

*pH of the 0.25% detergent + enzyme solution after 30' incubation

TABLE XII

Stability of different cellulases in 0.25% OMO® Total (or OMO® Neste pH 8.5).

		enzyme dosage	% of activity left					
preparation		% (ml)	pH*	0'	5'	10'	20'	30'
10	<u>Culture filtrates:</u>							
	ALKO4125	6	7.8	100	98	96	86	87
	ALKO4179	6	7.3	100	98	96	96	99
	ALKO4265	6	7.1	100	100	100	100	100
	ALKO4265	4	7.8	100	99	97	100	100
15	ALKO4237	4	7.8	100	100	100	99	100
	ALKO4237	2	7.3	100	99	97	99	99
	<u>Purified proteins from ALKO4237:</u>							
	20K-cellulase	1	8.2	100	100	99	93	100
20	50K-cellulase	1	7.8	100	95	92	95	94

*pH of the 0.25% detergent + enzyme solution after 30' incubation

TABLE XIII

Stability of different cellulases in 0.25% OMO® Color (pH 9.6–10)

		enzyme dosage	% of activity left					
	preparation	% (ml)	pH*	0'	5'	10'	20'	30'
30	<u>Culture filtrates:</u>							
	ALKO4125	6	9.6	100	(15)	(15)	(13)	(14)
	ALKO4179	6	8.3	100	97	100	97	99
	ALKO4265	6	9.1	100	100	100	100	100
	ALKO4265	4	8.5	100	93	95	99	98
	ALKO4237	4	8.5	100	98	96	96	99
	ALKO4237	2	9.1	100	93	95	99	98
<u>Purified proteins from ALKO4237:</u>								
40	20K-cellulase	1	9.8	100	99	100	100	100
	50K-cellulase	1	8.9	100	100	100	100	100

*pH of the 0.25% detergent + enzyme solution after 30' incubation

Example 14

Function of Neutral Cellulases in Detergents in HEC Substrate

The function of different neutral cellulases in detergents was determined by using hydroxyethylcellulose (HEC) as a substrate. The tested cellulase preparations were ALKO4265 and ALKO4237 concentrated culture filtrates and purified 20K- and 50K-cellulases from ALKO4237 strain. HEC substrates were prepared by dissolving 1% HEC into 0.25% detergent solutions. By using these substrates the activity against HEC (ECU/ml) was measured at 40° C. from each cellulase preparations as described in Example 1. Detergents and cellulase preparations used in these experiments are described in Example 13.

pH of the substrates:

HEC/buffer	pH 7
HEC/Colour Detergent Liquid	pH 7.5

-continued

pH of the substrates:	
HEC/OMO® Total	pH 7.8
HEC/OMO® Color	pH 9.7

TABLE XIV

ECU of the cellulase preparations in different detergents (compared as % from the ECU activity measured in pH 7 buffer)				
Activity %				
preparation	ECU/ buffer	ECU/ col.det.liquid	ECU/ OMO® Total	ECU/ OMO® Color
<u>culture filtrates:</u>				
ALKO4265	100	89	96	59
ALKO4237	100	97	95	40
<u>purified proteins:</u>				
20K-cellulase	100	100	93	81
50K-cellulase	100	92	79	46

ALKO4237 and ALKO4265 cellulase preparations and 20K- and 50K-cellulases function in all three tested detergents when using HEC as a substrate.

Example 15

Use of Neutral Cellulases in Detergents on Cotton Woven Fabrics

In this experiment is described the ability of the neutral cellulases to function as fabric-softening agent and to prevent furring and thus to reduce pilling tendency from cotton fabric after repeated launderings in detergents. The tested cellulase preparations were ALKO4237 concentrated culture filtrate and the purified 20K- and 50K-cellulases from ALKO4237 strain (Examples 1 and 9).

The washing experiment was carried out with a Launder-Ometer LP-2 (Atlas, Ill., USA). About 10 g of prewashed (Example 3) unbleached cotton woven fabric swatch was loaded into 1.2 liter container that contained 150 ml of 0.25% detergent solution with or without cellulase. Cellulase dosages were based on protein amounts. Detergent solutions were OMO® Total (Lever, UK) and Colour Detergent Liquid (Unilever, The Netherlands). A quantity of steel balls were added into each container to increase the mechanical action. The Launder-Ometer was run at 42 rpm for 0.5 or 1 hour at 40° C. The material was washed 4 times with intermediate rinsing and drying.

Weight loss (see Example 6) was used to describe the amount of fuzz removed from the fabrics surface.

TABLE XV

Weight loss of the fabrics after the first washing time with neutral cellulases in detergents.				
sample no	preparation	enzyme dosage as protein/ g fabric	time h	weight loss %
<u>In Colour Detergent Liquid:</u>				
1	—	—	1	0.05
2	ALKO4237	11	1	0.3

TABLE XV-continued

Weight loss of the fabrics after the first washing time with neutral cellulases in detergents.				
sample no	preparation	enzyme dosage as protein/ g fabric	time h	weight loss %
3	ALKO4237	22	1	0.7
4	20K-cellulase	2	1	0.1
5	20K-cellulase	5	1	0.5
6	20K-cellulase	8	1	1.0
7	50K-cellulase	2	1	0.1
8	50K-cellulase	5	1	0.2
9	—	—	0.5	0.2
10	20K-cellulase	8	0.5	0.5
<u>In OMO® Total:</u>				
11	—	—	1	0.03
12	20K-cellulase	8	1	1.1
13	—	—	0.5	0.1
14	20K-cellulase	8	0.5	0.7

In the Table XV it is shown that after the first washing in Launder-Ometer weight loss of the fabrics were increased clearly more with cellulase treated fabrics than with the fabrics treated with the sole detergent. Also weight loss was increased as a function of cellulase dosage and further with 20K-cellulase weight loss was increased when washing time was raised from 0.5 hour to 1 hour. 20K-cellulase worked equally well in Colour Detergent Liquid and in OMO® Total. These results indicate that particularly the 20K-cellulase and ALKO4237 cellulase preparation function in detergents as fuzz removing agents after already one wash time.

After three further washing times with samples 1, 2, 4 and 7 (Table XV) the evaluation of the fabrics was performed by a panel consisting of three persons. Panelists were asked to evaluate the softness and visual appearance of the treated fabrics as follows.

The softness of the fabrics:

- the fabric treated with cellulase is softer than the fabric treated without cellulase
- the fabric treated with cellulase is as soft as the fabric treated without cellulase
- the fabric treated with cellulase is harder than the fabric treated without cellulase

The results are shown in Table XVI.

Visual appearance of the fabrics was evaluated by ranking the fabrics on a score from 1 to 5. Score of 5 gave no few or pills and the fabric texture became more apparent. Score of 1 gave many pills and fuzz. Total score for each fabric was calculated and divided by the number of the panelists. The average score of the visual appearance of each fabric is shown in Table XVI.

TABLE XVI

Softness and visual appearance of the fabrics after 4 repeated washing times with neutral cellulases in detergents.

preparation	enzyme dosage as protein/ g fabric	time h	softness	visual appearance
In Colour Detergent Liquid:				
—	—	1		1
ALKO4237	11	1	100%: softer with cellulase	3.2
20K-cellulase	2	1	100%: softer with cellulase	3.7
50K-cellulase	2	1	100%: no difference	1.7

After the 4 treatments the cellulase treated fabrics had clearly better visual appearance than the fabrics that were treated with sole detergent. Thus fabrics treated with these cellulases maintained good appearance and the fuzziness was prevented after repeated washings compared to the fabric treated without cellulases. Also after 4 wash times the ALKO4237 and 20K-cellulase treated fabrics were softer than the fabric treated with sole detergent.

Green cotton fleecy knit swatches were washed at Launder-Ometer in Colour Liquid Detergent or in OMO® Total for 1 h 3 or 10 times with or without cellulases as described in Example 15.

The evaluation of the knits was performed by a panel consisting of three persons. Panelists were asked to evaluate the softness and visual appearance (both right and reverse sides) of the treated knits as described in Example 15. Weight loss of the knits was determined as described in Example 15. The results are shown in Table XVII.

After the 3 washing times the 2° K-cellulase treated knits had better visual appearance both on the right and reverse side than the knits treated with sole detergent. Knits treated 10 times with ALKO4237 cellulase preparation had clearly better visual appearance and brighter green colour than the knits treated only with detergent. The better visual appearance of the cellulase treated knits was detected already after 1 wash time (especially on the reverse side) and it was further developed during the additional washings. The cellulase treated knits were also softer than the knits treated with sole detergent.

TABLE XVII

Softness, weight loss and visual appearance of the fleecy knits after 3 or 10 repeated washing times with or without cellulases in detergents.
Before washings pH of the 0.25% Colour Detergent Solution was 7.9 and 8.4 of the 0.25% OMO® Total solution.

preparation	enzyme		pH	weight	softness	visual appearance	
	dosage as protein/g fabric	washing times	after washings	loss %		right	reverse side
Colour							
<u>Detergent Liquid</u>							
—	—	3	7.9	0.46		1	1
20K*	5	3	7.4	0.88	33%: softer with cellulase	1.5	2.7
—	—	10	8.0	1.46		1	1
A4237	20	10	7.9	2.80	100%: softer with cellulase	2.5	2.8
<u>OMO ® Total</u>							
—	—	10	8.3	0.57		1	1
A4237	20	10	8.2	1.57	100%: softer with cellulase	2.3	2.8

*= 20K-cellulase

Example 16

Use of Neutral Cellulases in Detergents on Cotton Fleecy Knit

In this experiment is described the ability of the neutral cellulases to function as fabric-softening agent and to prevent fuzzing and thus to reduce pilling tendency from coloured cotton fleecy knit after repeated launderings in detergents. The tested cellulase preparations were ALKO4237 concentrated culture filtrate and the purified 20K-cellulase from ALKO4237 strain (Examples 1 and 9).

Example 17

Use of Neutral Cellulases in Detergents on Aged Cotton Fleecy Knit

In this experiment is described the ability of the neutral cellulases to function as fabric-renewal and -softening agent.

Green cotton fleecy knit was washed 10 times, with intermediate drying, in Cylinda washing machine with programme 3 at 60° C., 10 ml of OMO® Color (Lever, UK). This was to simulate the washings of the knit in practice.

After 10 treatments this aged knit had unattractive and faded appearance with a lot of fuzz at the surface.

After these 10 repeated washes the fleecy knit was used for the washing experiments with or without cellulase. Knit swatches were washed at Launder-Ometer in Colour Liquid Detergent for 1 h 1 to 3 times as described in example 15 with intermediate rinsing and drying. The cellulase prepa-

rations used were ALKO4237 concentrated culture filtrate and purified 20K- and 50K-cellulases from ALKO4237 (Example 9).

The evaluation of the knits was performed by a panel consisting of three persons. Panelists were asked to evaluate the softness and visual appearance (both right and reverse sides) of the treated knits as described in Example 15. Weight loss of the knits was determined as described in Example 15. The results are shown in Table XVIII.

After one wash time ALKO4237 and 20K-cellulase treated knits had slightly better visual appearance than the knit treated with sole detergent. The good visual appearance and more attractive look was further developed to the 20K-cellulase treated knits after 2 and 3 wash times. Visual appearance was also improved after two wash times on the knits treated with 50K-cellulase compared to the knit treated with sole detergent. As general, the knits treated with cellulases had clearly improved and attractive look while the knits treated without cellulase had still unattractive and faded appearance.

calf intestine alkaline phosphatase. DNA ranging from 5–15 kb was isolated from an agarose gel using beta-agarase (Boehringer Mannheim, Germany) and used to construct the genomic ALKO4237 library.

The predigested Lambda DASH®II BamHI Vector Kit (Stratagene, USA) was used to construct the library and the instructions of the manufacturer were followed in all the subsequent steps. Briefly, about 200 ng of the size-fractionated DNA was ligated into 1 µg of DASH®II prepared arms, and packaged using Gigapack II packaging extract (Stratagene, USA). The titer of the library was determined by infecting *E. coli* XL 1-Blue MRA (P2)-cells with serial dilutions of the packaged phage and plating on NZY plates. The library was stored at 4° C. in SM-buffer, with 4% (v/v) chloroform. It was used for screening without amplification.

TABLE XVIII

Softness, weight loss and visual appearance of the aged fleecy knits after 1 to 3 repeated washing times with or without cellulases in detergents. Before washings pH of the 0.25% Color Detergent Solution was 7.9.

preparation	enzyme dosage as mg protein/g fabric	washing times	pH		weight loss %	softness	visual appearance	
			after washings				right side	reverse
—	—	1	ND		0		1	1
ALKO4237	20	1	ND		0.61	100%: no difference	1	1.5
20 K*	5	1	ND		0	100%: no difference	1.5	1.5
—	—	2	7.9		0.10		1	1
20 K*	5	2	7.7		0.46	100%: softer with cellulase	2.5	2.2
50 K*	5	2	7.7		0.26	100%: no difference	1	1.2
50 K*	15	2	7.3		0.49	100%: no difference	1	1.3
—	—	3	ND		0.31		1	1
20 K*	5	3	ND		0.88	100%: softer with cellulase	3.0	2.2

ND = not determined

* = 20 K- or 50 K-cellulase

Example 18

Isolation of the ALKO4237 Chromosomal DNA and Construction of the Genomic Library

Melanocarpus albomyces ALKO4237 was grown in shake flask cultures in potato dextrose (PD; Difco, USA)-medium at 42° C., 250 rpm for 3 days. The chromosomal DNA was isolated according to Raeder and Broda, *Lett. Appl. Microbiol.* 1:17–20 (1985). Briefly, the mycelium was washed with 20 mM EDTA and lysed in extraction buffer (200 mM Tris-HCl (pH 8.5), 250 mM NaCl, 25 mM EDTA, 0.5% SDS). The DNA was extracted with phenol and a mixture of chloroform:isoamyl alcohol (24:1 v/v). RNA was digested with RNase.

The chromosomal DNA was partially digested with Sau3A (Boehringer Mannheim, Germany) and treated with

Example 19

Amplification, Cloning and Sequencing of the 20K-cellulase DNA With Degenerate Primers

To amplify the 20K-cellulase gene by polymerase chain reaction (PCR), a pair of degenerate primers based on the peptide sequences (FIG. 17) (SEQ ID NOS:1–12) was synthesized. Primer 1 (429–32) (SEQ ID NO:38) was derived from the amino acids #8–14 of the N-terminal peptide #429 (FIG. 17) (SEQ ID NO:1), and primer 2 (fr28-16) (SEQ ID NO:39) was designed as the antisense strand for the amino acids #2–8 of the peptide fr28 (FIG. 17) (SEQ ID NO:11). Additional EcoRI restriction sites were added at the 5'-termini to facilitate the cloning of the amplified fragment.

Primer 1 (429–32) (SEQ ID NO:38)
EcoRI

5'- ATA GAATTC TA(C/T) TGG GA(C/T) TG(C/T) TG(C/T) AA(A/G) CC
Y W D C C K P

Primer 2(fr28-16) (SEQ ID NO:39)
EcoRI

that the about 35 kDa protein detected previously with the 20K-cellulase antiserum (Example 10) is a different gene product than the 20K-cellulase.

The 20K-cellulase of *Melanocarpus albomyces* appears to belong to family K of cellulases and family 45 of glycosyl hydrolases (Henrissat & Bairoch, *Biochem. J.* 293:781-788 (1993)). The 20K-cellulase shows homology (about 76% identify in 235 amino acid overlap) towards the *Humicola insolens* endoglucanase V (embl:a23635), but the 20K-cellulase has the surprising feature that it does not harbor the cellulose binding domain (CBD) and its linker, which are characteristic of the *Humicola insolens* endoglucanase V and other related endoglucanases (Schülein et al., 1993, In: Suominen & Reinikainen (eds), Foundation for Biotechnical and Industrial Fermentation Research, Helsinki, vol. 8, 109.; Saloheimo et al., 1994, *Mol. Microbiol.* 13, 219). This feature of the 20K-cellulase may account for the excellent performance of the enzyme in biostoning experiments (Example 10).

Example 21

Amplification, Cloning and Sequencing of 50K-cellulase DNA With Degenerate Primers

The peptides derived from the 50K-cellulase (Table IX) shared some homology towards *Humicola grisea* endoglucanase I (DDBJ:D63516). To amplify the 50 K-cellulase gene by polymerase chain reaction (PCR) a pair of degenerate primers based on the peptide sequences (Table IX) (SEQ ID NOS:13-24) was synthesized. Primer 1 (507-128) (SEQ ID NO:40) was derived from the amino acids #5-10 of the peptide #507 (Table IX)(SEQ ID NO:13), and primer 2 (509-rev) (SEQ ID NO:41) was designed as the antisense strand for the amino acids #4-9 of the peptide 509 (Table IX)(SEQ ID NO:14). The order of the two peptides in the protein—and the corresponding sense-antisense nature of the primers—was deduced from comparison with the *Humicola grisea* endoglucanase I.

Primer 1 (507-128) (SEQ ID NO:40)
5'-GA(C/T) GA(A/G) AC(A/C/G/T) GA(A/G) CA(C/T) (A/C)G
D E T E H R

Primer 2 (509-rev) (SEQ ID NO:41)
5'-TA (A/C/G/T)GC (A/C/G/T)CC (A/C/G/T)CC (A/C/G/T)GG (A/G)TT
Y A G G P N

In the PCR reaction, 1.5 µg of the purified ALKO4237 genomic DNA (Example 18) was used as the template. Dynazyme DNA polymerase (Finnzymes Ltd, Finland) was used according to the supplier's instructions.

Template DNA	(0.3 µg/µl)	5 µl
Primer 1	(0.5 µg/µl)	1 µl
Primer 2	(0.5 µg/µl)	1 µl
dNTPs	(2 mM)	5 µl
10xPCR buffer		10 µl
dH ₂ O		79 µl
Dynazyme	(2 U/µl)	1 µl
Total		102 µl

The PCR reaction was performed under the following conditions:

Step 1	95° C.	5 min
Step 2	95° C.	1 min
Step 3	56° C.	1 min
Step 4	72° C.	1 min
Step 5	go to "step 2"	29 more times
Step 6	72° C.	8 min
Step 7	4° C.	hold

Ten µl of reaction mixture was analyzed by agarose gel electrophoresis, and a single band corresponding to about 160 bp in length was detected. The remaining of the PCR product was loaded on a agarose gel electrophoresed, and the agarose section containing the DNA fragment was excised, and purified by the Magic PCR Preps (Promega, USA) method according to the supplier's instructions.

The isolated fragment was ligated with pBluescript II SK+(Stratagene, USA) plasmid which had been digested with EcoRV endonuclease, and ddT-tailed as described in Holton and Graham (1990) *Nucl. Acids Res.* 19, 1156. Competent *Escherichia coli* XL-Blue cells (Stratagene, USA) were transformed with the ligation mixture. Plasmid DNA from a few of the resulting colonies was isolated by the Magic Minipreps (Promega, USA) method according to the supplier's instructions. The plasmid DNA was analyzed by agarose electrophoresis, and one clone with expected characteristics was designated pALK1064.

The insert (161 bp) in pALK1064 was sequenced as described in Example 19, and was found to contain an ORF, which predicted a peptide homologous to *Humicola grisea* endoglucanase I (DDBJ:D63516). The ORF also encoded the peptide #612 (Table IX) (SEQ ID NO:21) from the purified 50K-cellulase. The PCR amplified DNA (in addition to the primers) corresponds to the nucleotides 404-530 in FIG. 21 (SEQ ID NO:32).

PCR with the primers 507 and 590-rev with ALKO4124 chromosomal DNA as template (Example 19) produced a

fragment of same size as from ALKO4237 DNA. This suggests that *Myriococcus* sp. ALKO4124 has a protein very similar to the 50K-cellulase of *Melanocarpus albomyces* ALKO4237. This is also supported by the fact that enzymes from both strains gave similar good results in biostoning experiments.

Example 22

Cloning and Sequencing the *Melanocarpus albomyces* ALKO4237 50K-Cellulase Gene

The genomic bank of *Melanocarpus albomyces* ALKO4237 was prepared for hybridization as described in Example 20. The purified PCR fragment carrying part of the 50K-cellulase gene (Example 21) was labeled with digoxigenin according to Boehringer, DIG DNA Labeling and Detection Nonradioactive, Application Manual. Hybridization was performed at 68° C. The positive clones were picked in SM buffer/chloroform, and purified with a second round of screening.

Under these conditions 10 positive clones were found. The large scale bacteriophage lambda DNA isolation from the clones was done according to Sambrook et al., 1989. The phage DNAs were analyzed by digestion of the DNA with several restriction enzymes, and the digested DNA was hybridized with the 50K-cellulase-specific PCR-probe. Four hybridizing fragments were isolated: about 2.8 kb SacI-XhoI fragment, about 5 kb SacI fragment, about 3.2 kb XhoI fragment, and about 2 kb EcoRI fragment. These were inserted into similarly cut pBluescript II SK+vector (Stratagene, USA), creating plasmids pALK1234, pALK1233, pALK1226 and pALK1227, respectively (FIG. 20).

The *Melanocarpus albomyces* ALKO4237 DNA was sequenced from the 50K-cellulase specific plasmids mentioned above. The sequencing protocol has been described in Example 19.

The DNA encoding the *Melanocarpus albomyces* 50K-cellulase is shown in FIG. 21 (A and B)(SEQ ID NO:32). The sequence reveals an ORF of about 1363 bp in length, interrupted by one intron. The ORF codes for 428 amino acids. The predicted protein has a molecular weight of 46.8 kDa and after signal peptide cleavage of 44.8 kDa. All the peptides in Table IX (SEQ ID NOS:13-14) are found in the predicted protein sequence (FIG. 21) (SEQ ID NO:33), although some amino acids identified with uncertainty during the peptide sequencing proved to be incorrect. The protein shows homology to *Humicola grisea* endoglucanase I (DDBJ:D63516).

Example 23

Amplification, Cloning and Sequencing of 50K-Cellulase B DNA With Degenerate Primers

The peptides derived from the 50K-cellulase B (Table X) (SEQ ID NOS:25-29) shared some homology towards *Humicola grisea* cellobiohydrolase I (DDBJ:D63515). To amplify the 50K-cellulase B gene by polymerase chain reaction (PCR) a pair of degenerate primers based on the peptide sequences (Table X) (SEQ ID NOS:25-29) was synthesized. Primer 1 (636) (SEQ ID NO:42) was derived from the amino acids #1-5 of the peptide #636 (Table X) (SEQ ID NO:29) (the first amino acid was guessed to be lysine, because this peptide was isolated after digestion with a protease cleaving after lysines), and primer 2 (534-rev) (SEQ ID NO:43) was designed as the antisense strand for the amino acids #3-8 of the peptide #534 (Table X)(SEQ ID NO:25). The order of the two peptides in the protein—and the corresponding sense-antisense nature of the primers—was deduced from comparison with the *Humicola grisea* cellobiohydrolase I.

Primer 1 (636) (SEQ ID NO:42)
5'-AA(A/G)CA(C/T)GA(A/G)TA(C/T)GG(A/C/G/T)AC
K H E Y G T

Primer 2 (534-rev) (SEQ ID NO:43)
5'-CC(A/G)TA(A/G)AA(A/G)TC(A/C/G/T)GG(A/G)TT
G Y F D P N

In the PCR reaction, 1.5 µg of the purified ALKO4237 genomic DNA (Example 18) was used as the template. Dynazyme DNA polymerase (Finnzymes Ltd, Finland) was used according to the supplier's instructions.

Template DNA	(0.3 µg/µl)	5 µl
Primer 1	(0.3 µg/µl)	1.7 µl
Primer 2	(0.3 µg/µl)	1.7 µl
dNTPs	(2 mM)	5 µl
10xPCR buffer		10 µl
dH ₂ O		80 µl
Dynazyme	(2 U/µl)	1 µl
Total		104.4 µl

The PCR reaction was performed under the following conditions:

Step 1	95° C.	5 min
Step 2	95° C.	1 min
Step 3	48° C.	1 min
Step 4	72° C.	2 min
Step 5	go to "step 2" 34 more times	
Step 6	72° C.	8 min
Step 7	4° C.	hold

Twenty µl of reaction mixture was analyzed by agarose gel electrophoresis, and a few bands were detected. One of the bands had an apparent size of 700 bp, which size was in agreement with size one would expect, when comparing with *Humicola grisea* cellobiohydrolase gene, particularly, if the fragment contained one or more introns. The PCR products were purified by the Magic PCR Preps (Promega, USA) method according to the supplier's instructions.

The isolated fragments was ligated with pBluescript II SK+(Stratagene, USA) plasmid which had been digested with EcoRV endonuclease, and ddT-tailed as described in Holton and Graham, *Nucl. Acids Res.* 19:1156 (1990). Competent *Escherichia coli* XL-Blue cells (Stratagene, USA) were transformed with the ligation mixture. Plasmid DNA from a few of the resulting colonies was isolated by the Magic Minipreps (Promega, USA) method according to the supplier's instructions. The plasmid DNA was analyzed by agarose electrophoresis, and one clone with about 700 bp insert was designated pALK1224.

The insert in pALK1224 was sequenced as described in Example 19, and was found to contain an ORF encoding the whole peptide #636 (SEQ ID NO:29) from the 50K-cellulase B (Table X). The ORF predicted a peptide homologous to *Humicola grisea* cellobiohydrolase I (DDBJ:D63515). The PCR amplified DNA (in addition to the primers) corresponds to the nucleotides 371-1023 in FIG. 23 (SEQ ID NO:39).

Example 24

Cloning and Sequencing the *Melanocarpus albomyces* ALKO4237 50K-Cellulase B Gene

The genomic bank of *Melanocarpus albomyces* ALKO4237 was prepared for hybridization as described in Example 20. The insert in pALK1224 was removed by digesting the plasmid with restriction endonucleases EcoRI and HindIII. The digested plasmid DNA was run by agarose electrophoresis. The agarose section containing the about 700 bp DNA fragment was excised, and purified by the Magic PCR Preps (Promega, USA) method according to the supplier's instructions.

The purified PCR fragment from pALK1224 carrying part of the 50K-cellulase B gene (Example 23) was labeled with

digoxigenin according to Boehringer, DIG DNA Labeling and Detection Nonradioactive, Application Manual. Hybridization was performed at 68° C. The positive clones were picked in SM buffer/chloroform, and purified with a second round of screening.

Under these conditions 3 positive clones were found. The large scale bacteriophage lambda DNA isolation from the clones was done according to Sambrook et al., in *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989. The phage DNAs were analyzed by digestion of the DNA with several restriction enzymes, and the digested DNA was hybridized with the 50K-cellulase B specific PCR probe. A hybridizing 3.5 kb NotI fragment was isolated, and inserted into similarly cut pBluescript II SK+vector (Stratagene, USA), creating plasmid pALK1229 (FIG. 22).

The extreme 5'-end of the gene was found by hybridizing the phage DNAs with 0.2 kb NotI-PstI-fragment from pALK1229. A hybridizing 2.4 kb PstI-fragment was isolated and inserted into similarly cut pBluescript II SK+ vector (Stratagene, USA), creating plasmid pALK1236 (FIG. 22).

Part of the inserts in pALK1229 and pALK1236 were sequenced as described in Example 19. The DNA encoding the *Melanocarpus albomyces* 50K-cellulase B is shown in FIG. 23(A and B)(SEQ ID NO:34). The sequence reveals an ORF of 1734 bp in length interrupted by five introns. The ORF codes for 452 amino acids. The predicted protein has a molecular weight of 49.9 kDa and after signal peptide cleavage of 47.6 kDa. All the peptides in Table X (SEQ ID NOS:25-29) are found in the predicted protein sequence (FIGS. 23A and B)(SEQ ID NO:35), although some amino acids identified with uncertainty during the peptide sequencing proved to be incorrect. The predicted protein shows homology to *Humicola grisea* cellobiohydrolase I (DDBJ:D63515) and other cellobiohydrolases. However, 50K-cellulase B has the surprising feature that it does not harbor the cellulose binding domain (CBD) and its linker, which is characteristic to *Humicola grisea* cellobiohydrolase I and many other cellobiohydrolases.

Example 25

Screening the *Melanocarpus albomyces* ALKO4237 Genomic Library with *Trichoderma reesei* Cellulases Genes

The genomic bank of *Melanocarpus albomyces* ALKO4237 was prepared for hybridization as described in Example 20.

A DNA fragment carrying *Trichoderma reesei* cbh1 specific DNA was isolated by cutting plasmid pTTc01 (FIG. 24) with restriction endonuclease HincII, and isolating the about 1.6 kb fragment from agarose gel after electrophoresis. A DNA fragment carrying *Trichoderma reesei* egl2 specific DNA was isolated by cutting plasmid pMS2 (FIG. 25) with restriction endonucleases BamHI and EcoRI, and isolating the about 1.5 kb fragment from agarose gel after electrophoresis. The cloning of the cbh1 gene is described in Teeri et al., *Bio/Technology* 1:696-699 (1983) and the DNA sequence is described in Shoemaker et al., *Bio/Technology* 1:691-696 (1983). The egl2 (originally called "egl3") gene is described in Saloheimo et al., *Gene* 63:11-21 (1988).

The fragments were labeled with digoxigenin according to Boehringer, DIG DNA Labeling and Detection Nonradioactive, Application Manual. Hybridization was performed at 68° C. with the cbh1 probe and at 60° C. with the egl2 probe. The positive clones were picked in SM buffer/chloroform, and purified with a second round of screening.

Under these conditions 13 cbh1 positive and 6 egl2 positive clones were found. One clone hybridized to both probes. The lambda DNA was isolated from the clones as described above. The phage DNAs were analyzed by digestion of the DNA with several restriction enzymes, and the digested DNA was hybridized with the cbh1 and egl2 probes. The clones were also hybridized with the 20K-cellulase-specific PCR fragment (Example 19). One clone (lambda-16) was clearly positive, and two other clones (lambda-8/1 and lambda-5/2) were weakly positive; all these clones were originally picked with the cbh1 probe.

An about 4 kb EcoRI fragment from lambda-16, which hybridized to both the *Trichoderma reesei* cbh1 probe and to the 20K-cellulase specific PCR fragment, was isolated from agarose gel after electrophoresis, and inserted into similarly cut pBluescript II SK+. The resulting plasmid was named pALK1230 (FIG. 26).

Part of the insert in pALK1230 was sequenced as described in Example 19. The DNA appears not to encode the 20K-cellulase, but codes for a protein homologous to several cellulases, particularly at the cellulose binding domain (CBD) area. Thus the gene product very likely has high affinity towards cellulosic material, and therefore this gene product was designated as protein-with-CBD. The sequence is shown in FIG. 27 (SEQ ID NO:36).

PCR reactions with the primers 636 (SEQ ID NO:42) and 534-rev (SEQ ID NO:43)(Example 23) were performed with the DNA from the 19 lambda clones as templates. One lambda clone, lambda-3, gave a band about 700 bp in size, similar to that in Example 23 when ALKO4237 chromosomal DNA was used as a template. This clone had originally been picked by the *Trichoderma* cbh1 probe. The lambda DNA was digested with several restriction endonucleases, and hybridized with the 50K-cellulase B specific probe. The clone showed similar restriction enzyme pattern as the 3 clones in Example 24. It is concluded that lambda-3 also carries the 50K-cellulase B gene.

Example 26

Fusion Proteins

A recombinant vector encoding the 20K-cellulase, 50K-cellulase or the 50K-cellulase B is prepared by fusing the cellulase encoding sequence with the sequence of *Trichoderma reesei* cellulase or hemicellulase or at least one functional domain of said cellulase or hemicellulase, as described in U.S. Pat. No. 5,298,405, WO 93/24621 and in Genbank submission L25310, incorporated herein by reference. Especially, the enzyme is selected from the group consisting of CBHI, CBHII, EGI, EGII, XYLI, XYLII and MANI, or a domain thereof, such as the secretion signal or the core sequence.

Fusion proteins can be constructed that contain an N-terminal mannanase or cellobiohydrolase or endoglucanase core domain or the core and the hinge domains from the same, fused to one of the *Melanocarpus* cellulase sequences. The result is a protein that contains an N-terminal mannanase or cellobiohydrolase or endoglucanase core or core and hinge regions, and a C-terminal *Melanocarpus* cellulase. The fusion protein contains both the *Trichoderma* mannanase or cellobiohydrolase or endoglucanase and the *Melanocarpus* cellulase activities of the various domains as provided in the fusion construct. Alternatively, mutations that modify the activities of the *Trichoderma* mannanase or cellobiohydrolase or endoglucanase, or the *Melanocarpus* cellulase activities, may be included in the constructions. In this case,

the fusion proteins contain both the modified *Trichoderma* enzyme activity and the *Melanocarpus* cellulase activity of the various domains as provided in the fusion construct.

Fusion proteins can also be constructed such that the mannanase or cellobiohydrolase or endoglucanase tail or a desired fragment thereof, is placed before one of the *Melanocarpus* cellulase sequences, especially so as to allow use of a nonspecific protease site in the tail as protease site for the recovery of the *Melanocarpus* cellulase part from the expressed fusion protein. Alternatively, fusion proteins can be constructed that provide for a protease site in a synthetic linker that is placed before one of the *Melanocarpus* cellulases, with or without the tail sequences.

Example 27

Hosts

The recombinant construct encoding the desired fusion proteins or *Melanocarpus* proteins are prepared as above, and transformed into a filamentous fungus such as *Aspergillus* spp., preferably *Trichoderma* spp.

Example 28

Trichoderma Background for 20K-Cellulase Production

In this example is described stone-washing experiments to determine the most suitable background of *Trichoderma* cellulases for 20K-cellulase production. The purpose of these experiments was to determine which *Trichoderma* cellulases would cause backstaining in stone-washing at neutral conditions.

Trichoderma reesei strain ALKO3620 (endoglucanase 2 gene is deleted) was chosen as host for these experiments. In previous studies *Trichoderma* EGII (endoglucanase II) enzyme has been shown to cause detrimental effects to cotton fibre structures and thus to weaken the strength properties of cotton-containing fabrics (In: Miettinen-Oinonen et al.: Effects of cellulases on cotton fiber and fabrics. In: *Proceedings of the TIWC96 Conference*, 1996, Vol. 1 (2), pp. 197).

Stone-washing experiments were performed at pH 6.5 and 7 as described in Example 3 except that no Berol was used.

The tested *Trichoderma* cellulase preparations were:

ALKO3133 (egl2 and cbh2 deleted)
ALKO3269 (egl2 and egl1 deleted)
ALKO3268 (egl2 and cbh1 deleted)

The dosage of *Trichoderma* preparations was about 2.5 mg (=low dosage, L) or about 5 mg (=high dosage, H) of total protein per g of fabric. 0.4 mg of purified 20K-cellulase per g of fabric was used when needed.

Results of color measurements of treated denim fabrics are shown in Table XIX.

The stone-washing results show that ALKO3269 (egl2 and egl1 deleted) background causes less backstaining at neutral conditions than ALKO3268 (egl2 and cbh1 deleted) or ALKO3133 (egl2 and cbh2 deleted) background. Thus the preferred host for 20K-cellulase production for biostoning is an ALKO3269-like strain. Although with higher 20K-cellulase concentrations the *Trichoderma* background has probably only very minor importance. An ALKO3269-like background is probably as good for 50K-cellulase and 50K-cellulase B production for biostoning as it is for 20K-cellulase production.

TABLE XIX

Color measurements of denim fabrics treated with different *Trichoderma* cellulase preparations with (+) or without (-) 20 K-cellulase.

preparation/ dosage	20 K +/-	Right side				Reverse side		
		pH	L	b	deltaE	L	b	deltaE
—	-	6.5	2.2	1.1	3.1	0.7	0.1	1.4
10 ALKO3620/L	-	6.5	2.2	2.6	3.0	-0.7	2.6	2.9
ALKO3620/L	+	6.5	5.5	4.0	7.7	-1.3	5.0	5.5
ALKO3133/L	-	6.5	1.9	2.2	3.7	0.2	1.6	2.3
ALKO3133/H	-	6.5	4.2	1.9	4.5	-1.5	3.3	4.8
ALKO3133/L	+	6.5	5.7	4.3	7.8	0.3	4.5	5.0
ALKO3133/H	+	6.5	8.5	4.0	9.4	-1.4	5.9	7.8
15 ALKO3269/L	-	6.5	2.9	1.9	4.4	0.8	0.8	1.6
ALKO3269/H	-	6.5	4.3	1.5	4.5	0.6	1.3	2.6
ALKO3269/L	+	6.5	6.6	4.2	8.7	1.1	4.0	4.3
ALKO3269/H	+	6.5	7.9	3.9	8.5	0.7	3.7	5.1
ALKO3268/L	-	6.5	2.9	1.7	3.7	0.1	1.8	3.0
ALKO3268/H	-	6.5	4.2	2.0	4.3	-0.7	3.4	5.0
ALKO3268/L	+	6.5	5.9	3.2	7.7	-1.2	4.5	6.0
20 ALKO3268/H	+	6.5	7.1	3.7	7.7	-2.0	5.8	7.3
—	-	7.0	2.9	0.8	2.6	0.7	0.5	1.5
ALKO3620/L	-	7.0	3.3	1.2	1.9	1.7	0.3	1.1
ALKO3620/L	+	7.0	6.7	3.4	5.6	1.1	3.2	2.9
ALKO3133/L	-	7.0	3.2	1.0	1.4	0.6	0.6	0.9
ALKO3133/L	+	7.0	5.9	3.7	5.5	0.1	4.3	3.1
25 ALKO3269/L	-	7.0	3.6	1.2	2.2	1.3	-0.3	1.3
ALKO3269/L	+	7.0	6.4	3.4	5.9	1.2	3.2	2.8
ALKO3268/L	-	7.0	2.9	1.4	3.9	0.5	0.4	2.5
ALKO3268/L	+	7.0	8.4	3.1	9.6	1.1	3.5	4.6

Example 29

Production of *Melanocarpus albomyces* ALKO423 7 20K-Cellulase in *T. reesei*

The *Trichoderma reesei* strains were constructed for *Melanocarpus albomyces* ALKO4237 20K-cellulase production. Strains produce *Melanocarpus* 20K-cellulase and are unable to produce *T. reesei*'s endoglucanase II and cellobiohydrolase I or endoglucanase I. Such preparations deficient in *Trichoderma* cellulolytic activity, and the making of same by recombinant DNA methods, are described in U.S. Pat. No. 5,298,405 or Suominen et al. (1993) High frequency one-step gene replacement in *Trichoderma reesei*. II. Effects of deletions of individual cellulase genes. *Mol. Gen. Genet.* 241: 523., incorporated herein by reference.

In construction of the *Melanocarpus albomyces* 20K-cellulase producing strains, the parental *Trichoderma reesei* strain ALKO3620 was transformed with the expression cassettes from the plasmid pALK1231 or pALK1235 (FIGS. 28 and 29). In the cassettes 20K-cellulase is expressed from the strong cbh1 promoter. The integration of the expression cassettes resulted in the replacements of the parental cbh1 (pALK1231) or the egl1 (pALK1235) genes.

In the host strain ALKO3620 the egl2 gene has been replaced by the 3.3 kb XbaI-BglII fragment of the ble gene from *Streptoalloteichus hindustanus* (Mattern et al. (1988) A vector of *Aspergillus* transformation conferring phleomycin resistance. *Fungal Genet. Newslett.* 35: 25.; Drocourt et al. (1990) Cassettes of the *Streptoalloteichus hindustanus* ble gene for transformation of lower and higher eukaryotes to phleomycin resistance. *Nucl. Acids Res.* 18: 4009.) using the recombinant DNA methods described in U.S. Pat. No. 5,298,405, incorporated herein by reference.

The plasmids pALK1231 and pALK1235 that were used in the construction of the *Melanocarpus* cellulase producing strains are identical to each other with respect to cbh1

promoter, 20K-cellulase gene and cbh1 terminator which are described below:

* *T. reesei* cbh1 (cellobiohydrolase 1) promoter: The promoter is from *Trichoderma reesei* VTT-D-80133 (Teeri et al. (1983) The molecular cloning of the major cellulase gene from *Trichoderma reesei*. *Bio/Technology* 1: 696.). The 2.2 kb EcoRI-SacII fragment (Karhunen et al. (1993) High frequency one-step gene replacement in *Trichoderma reesei*. I. Endoglucanase I overproduction. *Mol. Gen. Genet.* 241: 515.) was used in the construct. The sequence of the promoter area preceding the ATG was published by Shoemaker et al. (1983) Molecular cloning of exo-cellobiohydrolase from *Trichoderma reesei* strain L27. *Bio/Technology* 1. 691.). The last 15 nucleotides of the *T. reesei* L27 cbh1 promoter (the SacII site is underlined) are CCGCGGACTGGCATC (SEQ ID NO:44)(Shoemaker et al. 1983). The cbh1 promoter from the *T. reesei* strain VTT-D-80133 has been sequenced at Alko Research Laboratories, and an one nucleotide difference in the DNA sequence has been noticed within the above mentioned region. In the *T. reesei* strain VTT-D-80133 the sequence preceding the ATG is CCGCGGAACTG/C/GCATC (SEQ ID NO:45) (the SacII site is underlined, the additional cytosine in the DNA sequence is between the slashes).

The nucleotides missing from the promoter (10 bps after the SacII to the ATG) were added and the exact promoter fusion to the first ATG of the *Melanocarpus* 20K-cellulase (see below) was done by using the PCR (polymerase chain reaction) method. The fusion and the PCR fragment were sequenced to ensure that no errors had occurred in the reaction. In pALK1231 the promoter area is also functioning as a homologous DNA (together with the cbh1 3'-fragment; see below) to target the integration of the transforming DNA into the cbh1 locus.

Melanocarpus albomyces 20K-cellulase gene: The nucleotide sequence and deduced amino acid sequence of the 20K-cellulase gene encoding an 20 kDa cellulase is presented in Example 20 (FIG. 19)(SEQ ID NOS:30-31). A 0.9 kb fragment beginning from ATG-codon was used in both plasmids.

T. reesei cbh1 terminator: The 739 bp AvaII fragment (Karhunen et al. (1993) High frequency one-step gene replacement in *Trichoderma reesei*. I. Endoglucanase I overproduction. *Mol. Gen. Genet.* 241: 515.) starting 113 bp before the STOP codon of the cbh1 gene was added after the 20K-cellulase gene to ensure termination of transcription.

In addition the material described above the plasmid pALK1231 contains:

amdS gene: The gene has been isolated from *Aspergillus nidulans* VH1-TRSX6 and it is coding for acetamidase (Hynes et al. (1983) Isolation of genomic clones containing the amdS gene of *Aspergillus nidulans* and their use in the analysis of the structural and regulatory mutations. *Mol. Cell. Biol.* 3: 1430.). Acetamidase enables the strain to grow by using acetamide as the only nitrogen source and this characteristics has been used for selecting the transformants. The 3.1 kb fragment (SpeI-XbaI) from the plasmid p3SR2 (Kelly J. and Hynes M. (1985) Transformation of *Aspergillus niger* by the amdS gene of *Aspergillus nidulans*. *EMBO J.* 4: 475.) is used in the plasmids. The fragment contains 1007 bps of the promoter area, 1897 bps of the coding region (introns included) and the 183 bps terminator area of the amdS gene.

cbh1 3'-fragment: The fragment was isolated from *T. reesei* ALKO2466 by using plasmid rescue (1.7 kb, BamHI-EcoRI, starting 1.4 kb after the gene's STOP, Suominen et

al. (1993) High frequency one-step gene replacement in *Trichoderma reesei*. II. Effects of deletions of individual cellulase genes. *Mol. Gen. Genet.* 241: 523.). Strain ALKO2466 derives from the strain ALKO233 (Harkki et al. (1991) Genetic engineering of *Trichoderma* to produce strains with novel cellulase profiles. *Enzyme Microb. Technol.* 13: 227.). 3'-fragment is used together with the promoter area to target the 20K-cellulase gene to the cbh1 locus by homologous recombination.

The plasmid pALK1235 contains:

hph gene: The gene encoding HmB phosphotransferase is originally isolated from *E. coli* K-12 JM109 (Yanish-Perron et al. (1985) Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* 33: 103.) and it confers resistance to hygromycin B (HmB). Resistance to hygromycin (inactivated by phosphorylation by HmB phosphotransferase) was used for selecting the transformants. The hph gene together with the pki promoter and cbh2 terminator (see below) is isolated from plasmid pRLM_{ex}30 (Mach et al. (1994) Transformation of *Trichoderma reesei* based on hygromycin B resistance using homologous expression signals. *Curr. Genet.* 25: 567.) as a 2.2 kb NotI-PvuII fragment.

pki promoter: The about 0.75 kb pki (pyruvate kinase) promoter for expressing hph has been synthesized by PCR using *T. reesei* QM 9414 DNA as a template (Schindler et al. (1993) Characterization of the pyruvate kinase-encoding gene (pki1) of *Trichoderma reesei*. *Gene* 130: 271.).

cbh2 terminator: The cbh2 terminator sequence starts immediately after the STOP codon of the cbh2 gene (to the PvuII site 0.5 kb from the STOP codon; Mach et al. (1994) Transformation of *Trichoderma reesei* based on hygromycin B resistance using homologous expression signals. *Curr. Genet.* 25: 567.) and originates from plasmid pRLM_{ex}30.

egl15'-fragment: The 1.8 kb egl15'-fragment (ScaI-StuI) has been isolated from *T. reesei* QM 6a (Mandels and Reese (1957) Induction of cellulase in *Trichoderma viridae* as influenced by carbon sources and metals. *J. Bacteriol.* 73: 269.). This fragment is situated about 1.35 kb upstream from the egl1 coding region and it was used to target the integration of the the transforming DNA into the egl1 locus.

egl1 3'-fragment: The 1.6 kb egl1 3'-fragment (ScaI-XhoI) was, like the 5'-fragment, isolated from *T. reesei* QM 6a. The fragment is situated 0.3 kb downstream from the end of the egl1 gene and it was used for targeting of the transforming DNA into the egl1 locus.

The standard DNA methods described by Sambrook et al. (1989) In: *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. were used in construction of the vectors. The restriction enzymes, T4 DNA ligase, Klenow fragment of the DNA polymerase I, T4 DNA polymerase, polynucleotide kinase and Taq polymerase were from Boehringer Mannheim, Germany) and New England Biolabs (USA). Each enzyme was used according to the supplier's instructions. Plasmid DNA was isolated by using Qiagen columns (Qiagen GmbH, Germany) or Promega Magic Minipreps (Promega, USA) according to the manufacturer's protocols. The oligonucleotides used in the PCR-reactions and in sequencing reactions were synthesized by a ABI (Applied Biosystems, USA) 381A DNA Synthesizer. DNA sequencing was done as described in Example 19.

DNA fragments for cloning or transformations were isolated from low-melting-point agarose gels (FMC Bioproducts, USA) by β -agarase I treatment (New England Biolabs, USA) or by using the QIAEX Gel Extraction Kit (Qiagen GmbH, Germany) according to the supplier's instructions.

53

T. reesei ALKO3620 was transformed as described by Penttilä et al. (1987) A versatile transformation system for the cellulolytic filamentous fungus *Trichoderma reesei*. Gene 61: 155.) with the modifications described in Karhunen et al. (1993) High frequency one-step gene replacement in *Trichoderma reesei*. I. Endoglucanase I overproduction. *Mol. Gen. Genet.* 241: 515.). *T. reesei* transformants were transferred on a selective medium and purified through conidia. Transformants were stabilized by growing them on selective slants for two generations prior to sporulating on potato dextrose agar.

Example 30

Characteristics of the *Melanocarpus albomyces*
ALKO4237 20K-cellulase Producing Transformants

The purified transformants were grown in shake flasks in a medium containing 4% whey, 1.5% complex nitrogen source derived from grain, 5% KH₂PO₄ and 0.5% (NH₄)₂SO₄. Cultures were grown at 30° C. and 250 rpm for 7 days.

The culture supernatants were blotted directly onto nitrocellulose filters by a dot-blot apparatus. CBHI was detected by immunostaining using a CBHI specific monoclonal antibody CI-258 and EGI by specific monoclonal antibody EI-2 (Aho et al. (1991) Monoclonal antibodies against core and cellulose-binding domains of *Trichoderma reesei* cellobiohydrolases I and II and endoglucanase I. *Eur. J. Biochem.* 200: 643.) and the ProtoBlot Western blot AP system (Promega, USA) according to the recommendations of the manufacturer.

The *T. reesei* strains ALKO3620/pALK1231/14, ALKO3620/pALK1231/16, ALKO3620/pALK1231/20 and ALKO3620/pALK1231/59 do not contain the *cbh1* gene. The *cbh1* gene is replaced by the *amdS* marker gene and the 20K-cellulase construct in pALK1231 expression cassette. The *cbh1* gene replacement was verified in Southern hybridisations. The *T. reesei* strains ALKO3620/pALK1235/40 and ALKO3620/pALK1235/49 do not contain the *egl1* gene. The *egl1* gene is replaced by the *hph* marker gene and the 20K-cellulase construct in pALK1235 expression cassette. The *egl1* gene replacement was verified in Southern hybridisations. The host strain ALKO3620 used in the transformations is deficient of the *egl2* gene (replaced by *ble* gene from *Streptoalloteichus hindustanus* (Mattern et al., 1988, Drocourt et al., 1990). Thus the strains do not produce *Trichoderma*'s cellulase components EGII and CBHI or EGI.

Samples from the culture supernatants were run on polyacrylamide slab gels containing 0.1% SDS on Bio-Rad Mini Protean II electrophoresis system (USA). The polyclonal antibody prepared against the purified 20K-cellulase was used to detect the produced protein in Western blots. In the detection, Promega's ProtoBlot® AP System was used. The Western result is shown in FIG. 30. The transformants ALKO3620/pALK1235/49, ALKO3620/pALK1235/40, ALKO3620/pALK1231/14 and ALKO3620/pALK1231/16 (lanes 1, 2, 4 and 5) produce a protein which reacts with the polyclonal 20K-cellulase antiserum. The size of the protein produced by transformants is same as the size of purified 20K-cellulase (lane 6). ALKO3620 (lane 3) does not produce corresponding protein.

The endoglucanase activities of the transformants were determined as described in Example 10. When 2% carboxymethylcellulose (CMC) was used as a substrate reaction temperature was lifted up to 70° C. and thus the endoglucanase activity of ALKO3620 was heat inactivated.

54

When using 1% hydroxyethylcellulose as a substrate heat inactivation was performed before enzymatic activity measurements. Samples from growth medium were diluted to 0.05 M HEPES, pH 7.0-buffer and incubated 20 min in 70° C. Heat inactivation of endoglucanase I (the major endoglucanase left in ALKO3620) was almost complete. The activity of *egl1*-negative transformants dropped about 30% in heat inactivation which indicates the minor heat inactivation of 20K-cellulase. The endoglucanase activities are presented in Table XX. When HEC was the substrate, the 20K-cellulase activity was extrapolated to the activity before the heat treatment by dividing the activity obtained after the heat treatment with 0.7.

Table XX

The endoglucanase activities of *T. reesei* transformants producing
Melanocarpus albomyces 20 K-cellulase.

Substrate	CMC 70° C., pH 7.0	20 K-cellulase activity (artificial units/ml)
		HEC 50° C., pH 7.0
ALKO4237	—*	100**
ALKO3620	50***	38***
ALKO3620/pALK1231/14	2400	350
ALKO3620/pALK1231/16	2600	350
ALKO3620/pALK1231/20	6500	750
ALKO3620/pALK1231/59	6800	750
ALKO3620/pALK1235/40	2400	325
ALKO3620/pALK1235/49	2100	350

*not measured

**not heat inactivated, contains also 50 K-cellulase, 50 K-cellulase B and other cellulase activities.

***activity due to *Trichoderma cellulases*

The endoglucanase activities of the *T. reesei* host strain ALKO3620 are almost totally heat inactivated at 70° C. *Melanocarpus albomyces* 20K-cellulase producing transformants produce substantial amounts of relative heat stable 20K-cellulase. The endoglucanase production level of transformants is several times higher than that of 20K-cellulase parental strain ALKO4237.

Example 31

Production of *Melanocarpus albomyces*
ALKO423750K-Cellulase in *T. reesei*

The *Trichoderma reesei* strains were constructed for *Melanocarpus albomyces* ALKO4237 50K-cellulase production. Strains produce *Melanocarpus* 50K-cellulase and are unable to produce *T. reesei*'s endoglucanase II and cellobiohydrolase I or endoglucanase I. In construction of the *Melanocarpus albomyces* 50K-cellulase producing strains, the parental *Trichoderma reesei* strain ALKO3620 was transformed with the expression cassettes from the plasmid pALK1238 or pALK1240 (FIGS. 31 and 32). In the cassettes 50K-cellulase is expressed from the strong *cbh1* promoter. The integration of the expression cassettes results in the replacements of the parental *cbh1* (pALK1238) or the *egl1* (pALK1240) genes. Cloning and transformation were done as described in Example 29, except that 20K-cellulase gene was replaced by 50K-cellulase gene (1.7 kb fragment beginning from ATG-codon) described in Example 22. The *Melanocarpus albomyces* 50K-cellulase producing transformants are then characterized similar to example 30 with modifications obvious to a person skilled in the art. The *Melanocarpus albomyces* 50K-cellulase B and protein-with-

CBD producing transformants can be created similar to Examples 29 and 30 with modifications obvious to a person skilled in the art.

Having now fully described the invention, it will be understood by those with skill in the art that the invention

may be performed within a wide and equivalent range of conditions, parameters and the like, without affecting the spirit or scope of the invention or any embodiment thereof. All references cited herein are fully incorporated herein by reference.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 45

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..30
- (D) OTHER INFORMATION:/label= No_429

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ala	Asn	Gly	Gln	Ser	Thr	Arg	Tyr	Trp	Asp	Cys	Cys	Lys	Pro	Ser	Cys
1				5					10					15	
Gly	Trp	Arg	Gly	Lys	Gly	Pro	Val	Asn	Gln	Pro	Val	Tyr	Ser		
			20					25					30		

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..7
- (D) OTHER INFORMATION:/label= No_430

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Tyr	Gly	Gly	Ile	Ser	Ser	Arg
1				5		

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

-continued

(vi) ORIGINAL SOURCE:
(A) ORGANISM: *Melanocarpus albomyces*
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION:1..4
(D) OTHER INFORMATION:/label= No_431

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Cys Gly Trp Arg

1

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: *Melanocarpus albomyces*
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION:1..6
(D) OTHER INFORMATION:/label= No_432

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Pro Ser Cys Gly Trp Arg

1

5

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: *Melanocarpus albomyces*
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION:1..6
(D) OTHER INFORMATION:/label= No_433

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Tyr Trp Asp Cys Cys Lys

1

5

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: *Melanocarpus albomyces*
(B) STRAIN: ALKO4237

-continued

(ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION:1..17
 (D) OTHER INFORMATION:/label= No_439

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Gln Glu Cys Asp Ser Phe Pro Glu Pro Leu Lys Pro Gly Cys Gln Trp
1 5 10 15

Arg

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 8 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Melanocarpus albomyces
 (B) STRAIN: ALKO4237

(ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION:1..8
 (D) OTHER INFORMATION:/label= fr9

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Arg His Asp Asp Gly Gly Phe Ala
1 5

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Melanocarpus albomyces
 (B) STRAIN: ALKO4237

(ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION:1..7
 (D) OTHER INFORMATION:/label= fr14

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Tyr Trp Asp Cys Cys Lys Pro
1 5

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Melanocarpus albomyces
 (B) STRAIN: ALKO4237

(ix) FEATURE:
 (A) NAME/KEY: Peptide

-continued

(B) LOCATION:1..18
(D) OTHER INFORMATION:/label= fr16

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Gly	Lys	Gly	Pro	Val	Asn	Gln	Pro	Val	Tyr	Ser	Cys	Asp	Ala	Asn	Phe
1				5					10					15	

Gln Arg

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Melanocarpus albomyces
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION:1..10
(D) OTHER INFORMATION:/label= fr17

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Val	Gln	Cys	Pro	Glu	Glu	Leu	Val	Ala	Arg
1			5						10

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Melanocarpus albomyces
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION:1..15
(D) OTHER INFORMATION:/label= fr28

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Asp	Trp	Phe	Gln	Asn	Ala	Asp	Asn	Pro	Ser	Phe	Thr	Phe	Glu	Arg
1				5					10				15	

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Melanocarpus albomyces
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION:1..30
(D) OTHER INFORMATION:/label= fr30

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Thr Met Val Val Gln Ser Thr Ser Thr Gly Gly Asp Leu Gly Ser Asn
1 5 10 15
His Phe Asp Leu Asn Ile Pro Gly Gly Val Gly Leu Phe
 20 25 30

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..10
- (D) OTHER INFORMATION:/label= No_507

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Val Tyr Leu Leu Asp Glu Thr Glu His Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..12
- (D) OTHER INFORMATION:/label= No_509

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Xaa Xaa Leu Asn Pro Gly Gly Ala Tyr Tyr Gly Thr
1 5 10

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..17
- (D) OTHER INFORMATION:/label= No_563

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Met Ser Glu Gly Ala Glu Cys Glu Tyr Asp Gly Val Cys Asp Lys Asp
1 5 10 15

Gly

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..14
- (D) OTHER INFORMATION: /label= No_565

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Asn Pro Tyr Arg Val Xaa Ile Thr Asp Tyr Tyr Gly Asn Ser
1 5 10

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /label= No_603

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Asp Pro Thr Gly Ala Arg Ser Glu Leu Asn Pro Gly Gly Ala Tyr Tyr
1 5 10 15

Gly Thr Gly Tyr Xaa Asp Ala Gln
20

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..13
- (D) OTHER INFORMATION: /label= No_605

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Xaa Xaa Val Pro Asp Tyr His Gln His Gly Val Asp Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..12
- (D) OTHER INFORMATION:/label= No_610

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Asn Glu Met Asp Ile Xaa Glu Ala Asn Ser Arg Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..25
- (D) OTHER INFORMATION:/label= No_611

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Leu Pro Xaa Gly Met Asn Ser Ala Leu Tyr Leu Ser Glu Met Asp Pro
1 5 10 15

Thr Gly Ala Arg Ser Glu Leu Asn Pro
20 25

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..21
- (D) OTHER INFORMATION:/label= No_612

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

Val Glu Pro Ser Pro Glu Val Thr Tyr Ser Asn Leu Arg Xaa Gly Glu
1 5 10 15
Ile Xaa Gly Xaa Phe
 20

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..19
- (D) OTHER INFORMATION:/label= No_619

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Asp Gly Cys Gly Trp Asn Pro Tyr Arg Val Val Ile Thr Thr Asp Tyr
1 5 10 15
Tyr Asn Asn

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..10
- (D) OTHER INFORMATION:/label= No_620

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

Leu Pro Cys Gly Met Xaa Ser Ala Leu Tyr
1 5 10

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION:1..22
- (D) OTHER INFORMATION:/label= No_621

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Ala Asp Gly Cys Gln Pro Arg Thr Asn Tyr Ile Val Leu Asp Asp Leu
1 5 10 15
Leu His Pro Xaa Xaa Gln
 20

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..9
- (D) OTHER INFORMATION: /label= No_534

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

Val Gly Asn Pro Asp Phe Tyr Gly Lys
1 5

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..8
- (D) OTHER INFORMATION: /label= No_535

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Phe Gly Pro Ile Gly Ser Thr Tyr
1 5

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Melanocarpus albomyces
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..12
- (D) OTHER INFORMATION: /label= No_631

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

Leu Ser Gln Tyr Phe Ile Gln Asp Gly Glu Arg Lys
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..11
- (D) OTHER INFORMATION: /label= No_632

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Phe Thr Val Val Ser Arg Phe Glu Glu Asn Lys
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..19
- (D) OTHER INFORMATION: /label= No_636

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

His Glu Tyr Gly Thr Asn Val Gly Ser Arg Phe Tyr Leu Met Asn Gly
 1 5 10 15

Pro Asp Lys

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 936 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 33..115
- (D) OTHER INFORMATION: /codon_start= 33
/product= "20K-cellulase"

(ix) FEATURE:

-continued

(A) NAME/KEY: exon
(B) LOCATION:187..435
(D) OTHER INFORMATION:/product= "20K-cellulase"

(ix) FEATURE:
(A) NAME/KEY: exon
(B) LOCATION:506..881
(D) OTHER INFORMATION:/product= "20K-cellulase"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

TCGCCCTAA CCGAGAACCA AAGACTCCAA GAATGCGCTC TACTCCCGTT CTCGCGCCCC	60
TCCTGGCGCG AGCATTGCCC CTCGGGGCCC TCGCCGCCAA CGGTCACTCC ACGAGGTAAC	120
TGATCACCCG CCTCATTACG CGTGCCGACC GGACCGCGTT CAGGGCTCAC TGCTACCCGC	180
ATCCAGATAC TGGGACTGCT GCAAGCCGTC GTGCGGCTGG CGCGGAAAGG GCCCCGTGAA	240
CCAGCCCCTC TACTCGTGCG ACGCCAACCT CCAGCGCATC CAGACTTCG ATGCCGTCTC	300
GGGCTGCGAG GCGGGCCCCG CCTTCTCGTG CGCCGACCAC AGCCCTGGG CCATTAATGA	360
CAACCTCTCG TACGGCTTCG CGGCGACTGC ACTCAGCGGC CAGACCGAGG AGTCGTGGTG	420
CTGTGCTCGT TACGCGTGAG TGTGCTGGG CCAACGTCG GTGATTCCGG AGTTCAGACC	480
ACTGACCCAG CGACCCGCTC GCCAGTCTGA CCTTACATC GGGTCCCGTG GCCGGCAAGA	540
CCATGGTCGT CCAGTCGACC AGCACGGGCG GCGACCTCGG CAGCAACAC TTCGACCTCA	600
ACATCCCGG CGGCGCGCTC GGCTCTTGG ACGGCTGCAC TCCCAGTTC GCGGGCTCC	660
CGGGCGCACG GTACGGCGGC ATCTCGTCGC GCCAGGAGTG CGACTCGTTC CCCGAGCCGC	720
TCAAGCCCG CTGCCAGTGG CGCTTCGACT GGTTCAGAA CGCCGACAAC CCGTCCTTTA	780
CCTTCGAGCG GGTCCAGTGC CCCGAGGAGC TGGTCGCTCG GACCGGTGC AGGCGCCACG	840
ACGACGGCGG CTTCGCCGTC TTCAAGGCCC CCAGCGCCTG ATCCGTTTTT GGCAGTGTG	900
CGTGTGACGG CAGCTACGTG GAACGACCTG GAGCTC	936

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 235 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Melanocarpus albomyces
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: Protein
(B) LOCATION:1..235
(D) OTHER INFORMATION:/label= 20K-cellulase

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

Met Arg Ser Thr Pro Val Leu Arg Ala Leu Leu Ala Ala Leu Pro	
1	5 10 15
Leu Gly Ala Leu Ala Ala Asn Gly Gln Ser Thr Arg Tyr Trp Asp Cys	
20	25 30
Cys Lys Pro Ser Cys Gly Trp Arg Gly Lys Gly Pro Val Asn Gln Pro	
35	40 45
Val Tyr Ser Cys Asp Ala Asn Phe Gln Arg Ile His Asp Phe Asp Ala	
50	55 60
Val Ser Gly Cys Glu Gly Gly Pro Ala Phe Ser Cys Ala Asp His Ser	
65	70 75 80

-continued

Pro Trp Ala Ile Asn Asp Asn Leu Ser Tyr Gly Phe Ala Ala Thr Ala
85 90 95

Leu Ser Gly Gln Thr Glu Glu Ser Trp Cys Cys Ala Cys Tyr Ala Leu
100 105 110

Thr Phe Thr Ser Gly Pro Val Ala Gly Lys Thr Met Val Val Gln Ser
115 120 125

Thr Ser Thr Gly Gly Asp Leu Gly Ser Asn His Phe Asp Leu Asn Ile
130 135 140

Pro Gly Gly Gly Val Gly Leu Phe Asp Gly Cys Thr Pro Gln Phe Gly
145 150 155 160

Gly Leu Pro Gly Ala Arg Tyr Gly Gly Ile Ser Ser Arg Gln Glu Cys
165 170 175

Asp Ser Phe Pro Glu Pro Leu Lys Pro Gly Cys Gln Trp Arg Phe Asp
180 185 190

Trp Phe Gln Asn Ala Asp Asn Pro Ser Phe Thr Phe Glu Arg Val Gln
195 200 205

Cys Pro Glu Glu Leu Val Ala Arg Thr Gly Cys Arg Arg His Asp Asp
210 215 220

Gly Gly Phe Ala Val Phe Lys Ala Pro Ser Ala
225 230 235

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1894 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
(B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: exon
(B) LOCATION: 233..838
(D) OTHER INFORMATION: /product= "50K-cellulase"

(ix) FEATURE:

- (A) NAME/KEY: exon
(B) LOCATION: 916..1596
(D) OTHER INFORMATION: /product= "50K-cellulase"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

GAATTCGGGG GTTGCCAGGG AGTCGTACAG GGGTGGGTGG AGGGGGATGG GGGATGGAAG 60

GGGGATGGAG AAGAAAGCAT ATATGGGACG TTTGTGCTCG CCGGCTCCCC TCTGCCACGT 120

TCCCTTGCCT CCTTGCCTGG GTTGTGTGTTG GTCTTCCCTT CACCATCCGA CAAACCAACC 180

TGCTGCGGGT GAACTCGCAG AGCGCCTTCG GACGACGACA GACAGACGCA CCATGACTCG 240

CAACATCGCC CTGCTCGGCG CCGCGTCGGC GCTCCTGGGC CTCGCCACG GCCAGAAGCC 300

GGGCGAGACG CCCGAGGTGC ACCCGCAGCT GACGACGTTT CCGTGCACCA AGGCGGACGG 360

GTGCCAGCCG CGGACCAACT ACATTGTGCT GGACTCGCTG TCGACCCCGG TGCACCAGGT 420

GGACAACGAC TACAACGCG GCGACTGGGG GCAGAAGCCC AACGCGACGG CGTGCCCGGA 480

CCTCGAGTCG TCGCGCGCA ACTGCATCAT GGAGGCGGTG CCCGACTACA GCCAGCACGG 540

CGTCACGACG AGCGACACGT CGCTGCGCCT GCAGCAGCTC GTCGACGGCC GCCTCGTCAC 600

GCCGCGGCTC TACCTGCTCG ACGAGACCGA GCACCGCTAC GAGATGATGC ACCTGACCGG 660

-continued

```

CCAGGAGTTC ACCTTTGAGG TCGACGCCAC CAAGCTGCCC TCGGGCATGA ACAGCGCCCT 720
CTACCTGTCC GAGATGGACC CGACCGGCGC CCGGAGCGAG CTCACCCCG GCGGTGCCTA 780
CTACGGCACC GGCTACTGCG ACGCCAGTG CTTCGTGACG CCATTCATCA ACGGCATTGT 840
GAGTGTTCCT CTTTGGCCCC CCCCTGAAA ATAGATGTAC CTGGGTGCTA ACCCGGGGT 900
GTGCGACCAA AACAGGGCAA CATCGAGGGC AAGGGCTCGT GCTGCAACGA GATGGACATC 960
TGGGAGGCCA ACTCGCGGGC GACGCACGTG GCGCCGACA CGTGCAACCA GACGGGTCTG 1020
TACATGTGCG AGGGCCCGCA GTGCGAGTAC GACGGCGTGT GCGACAAGGA CGGGTGCGGG 1080
TGGAACCCGT ACCGGGTCAA CATCACCGAC TACTACGGCA ACTCGGACGC GTTCCGCGTC 1140
GACACGCGGC GGCCCTTCAC CGTGGTGACG CAGTTCCCGG CCGACGCCGA GGGCCGGCTC 1200
GAGAGCATCC ACCGGCTGTA CGTGCAGGAC GGCAAGGTGA TCGAGTCGTA CGTCGTGAC 1260
GCGCCGGGCC TGCCCGGAC GACTCGCTC AACGACGAGT TCTGCGCCGC CACGGGCGCC 1320
GCGCGCTACC TCGACCTCGG CGGCACCGCG GGCATGGGCG ACGCCATGAC GCGCGGCATG 1380
GTGCTGGCCA TGAGCATCTG GTGGGACGAG TCCGGCTTCA TGAAGTGGCT CGACAGCGGC 1440
GAGGCCGGCC CCTGCCTGCC CGACGAGGGC GACCCCAAGA ACATTGTCAA GGTGAGGCC 1500
AGCCCCGAGG TCACCTACAG CAACCTGCGC TGGGGCGAGA TCGGGTCGAC CTTTGAGGCC 1560
GAGTCCGACG ACGACGCGCA CGGCGACGAC TGCTAGATAA CTAAGTAGTG GCGGAAAGG 1620
GCGGGGATG CGTAACTTAC ATACAGCCCG GAGTTGTTTT GAGTGTAGAG TATTGAGCTT 1680
TCGATGTGTT AGTTGAGTGG AATGAAAAAT TCGCGTCTTT GCCCGGTTGG TTGCGATAAA 1740
CAATAGTCGG CTGGTGCAAT TGTGACACTT CAATTGCGCT GTTGCTTGG TGACAGACAC 1800
GGCAGCGTCG ATGACCCGAC ACCCAGAATA ATTCGCATGG TTGATTATGT TATTGTGCTT 1860
TAAATCGGAG GCTGATGCTC ATCTCTTCGA ATTC 1894

```

(2) INFORMATION FOR SEQ ID NO: 33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 428 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALK04237

(ix) FEATURE:

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..428
- (D) OTHER INFORMATION: /label= 50K-cellulase

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

```

Met Thr Arg Asn Ile Ala Leu Leu Gly Ala Ala Ser Ala Leu Leu Gly
1           5           10           15
Leu Ala His Gly Gln Lys Pro Gly Glu Thr Pro Glu Val His Pro Gln
20           25           30
Leu Thr Thr Phe Arg Cys Thr Lys Ala Asp Gly Cys Gln Pro Arg Thr
35           40           45
Asn Tyr Ile Val Leu Asp Ser Leu Ser His Pro Val His Gln Val Asp
50           55           60
Asn Asp Tyr Asn Cys Gly Asp Trp Gly Gln Lys Pro Asn Ala Thr Ala
65           70           75           80

```

-continued

Cys Pro Asp Val Glu Ser Cys Ala Arg Asn Cys Ile Met Glu Gly Val
 85 90 95
 Pro Asp Tyr Ser Gln His Gly Val Thr Thr Ser Asp Thr Ser Leu Arg
 100 105 110
 Leu Gln Gln Leu Val Asp Gly Arg Leu Val Thr Pro Arg Val Tyr Leu
 115 120 125
 Leu Asp Glu Thr Glu His Arg Tyr Glu Met Met His Leu Thr Gly Gln
 130 135 140
 Glu Phe Thr Phe Glu Val Asp Ala Thr Lys Leu Pro Cys Gly Met Asn
 145 150 155 160
 Ser Ala Leu Tyr Leu Ser Glu Met Asp Pro Thr Gly Ala Arg Ser Glu
 165 170 175
 Leu Asn Pro Gly Gly Ala Tyr Tyr Gly Thr Gly Tyr Cys Asp Ala Gln
 180 185 190
 Cys Phe Val Thr Pro Phe Ile Asn Gly Ile Gly Asn Ile Glu Gly Lys
 195 200 205
 Gly Ser Cys Cys Asn Glu Met Asp Ile Trp Glu Ala Asn Ser Arg Ala
 210 215 220
 Thr His Val Ala Pro His Thr Cys Asn Gln Thr Gly Leu Tyr Met Cys
 225 230 235 240
 Glu Gly Ala Glu Cys Glu Tyr Asp Gly Val Cys Asp Lys Asp Gly Cys
 245 250 255
 Gly Trp Asn Pro Tyr Arg Val Asn Ile Thr Asp Tyr Tyr Gly Asn Ser
 260 265 270
 Asp Ala Phe Arg Val Asp Thr Arg Arg Pro Phe Thr Val Val Thr Gln
 275 280 285
 Phe Pro Ala Asp Ala Glu Gly Arg Leu Glu Ser Ile His Arg Leu Tyr
 290 295 300
 Val Gln Asp Gly Lys Val Ile Glu Ser Tyr Val Val Asp Ala Pro Gly
 305 310 315 320
 Leu Pro Arg Thr Asp Ser Leu Asn Asp Glu Phe Cys Ala Ala Thr Gly
 325 330 335
 Ala Ala Arg Tyr Leu Asp Leu Gly Gly Thr Ala Gly Met Gly Asp Ala
 340 345 350
 Met Thr Arg Gly Met Val Leu Ala Met Ser Ile Trp Trp Asp Glu Ser
 355 360 365
 Gly Phe Met Asn Trp Leu Asp Ser Gly Glu Ala Gly Pro Cys Leu Pro
 370 375 380
 Asp Glu Gly Asp Pro Lys Asn Ile Val Lys Val Glu Pro Ser Pro Glu
 385 390 395 400
 Val Thr Tyr Ser Asn Leu Arg Trp Gly Glu Ile Gly Ser Thr Phe Glu
 405 410 415
 Ala Glu Ser Asp Asp Asp Gly Asp Gly Asp Asp Cys
 420 425

(2) INFORMATION FOR SEQ ID NO: 34:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2000 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

-continued

(A) ORGANISM: *Melanocarpus albomyces*
(B) STRAIN: ALKO4237

(ix) FEATURE:
(A) NAME/KEY: exon
(B) LOCATION:154..729
(D) OTHER INFORMATION:/product= "50K-cellulase B"

(ix) FEATURE:
(A) NAME/KEY: exon
(B) LOCATION:810..946
(D) OTHER INFORMATION:/product= "50K-cellulase B"

(ix) FEATURE:
(A) NAME/KEY: exon
(B) LOCATION:1018..1230
(D) OTHER INFORMATION:/product= "50K-cellulase B"

(ix) FEATURE:
(A) NAME/KEY: exon
(B) LOCATION:1308..1551
(D) OTHER INFORMATION:/product= "50K-cellulase B"

(ix) FEATURE:
(A) NAME/KEY: exon
(B) LOCATION:1637..1767
(D) OTHER INFORMATION:/product= "50K-cellulase B"

(ix) FEATURE:
(A) NAME/KEY: exon
(B) LOCATION:1831..1888
(D) OTHER INFORMATION:/product= "50K-cellulase B"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

CCCGGTCTGG AGACGGGGAG CGGCCAGCG ACGCAGGATA AGAAGGCGAC GACCGCGCCT	60
CCGAGCCAGG CCCAGGACAG CAGGAGAAGT CGCCACGCGC AAGCAGCACG CCCGATCGAC	120
AGTGTCCTCG TCTGCCACA GCACTCTGCA ACCATGATGA TGAAGCAGTA CCTCCAGTAC	180
CTCGCGGCGG CGCTGCCGCT CGTCGGCCTC GCCGCCGGCC AGCGCGCTGG TAACGAGACG	240
CCCGAGAACC ACCCCCCGCT CACCTGGCAG AGGTGCACGG CCCCAGGCAA CTGCCAGACC	300
GTGAACGCCG AGGTGTCAT TGACGCCAAC TGGCGCTGGC TGCACGACGA CAACATGCAG	360
AACTGCTACG ACGGCAACCA GTGGACCAAC GCCTGCAGCA CCGCCACCGA CTGCGCTGAG	420
AAGTGCATGA TCGAGGGTGC CGCGGACTAC CTGGGCACCT ACGGCGCCTC GACCAGCGGC	480
GACGCCCTGA CGCTCAAGTT CGTCACCAAG CACGAGTACG GCACCAACGT CGGCTCGCGC	540
TTCTACCTCA TGAACGGCCC GGACAAGTAC CAGATGTTCA ACCTCATGGG CAACGAGCTT	600
GCCTTTGACG TCGACCTCTC GACCGTCGAG TCGGCATCA ACAGCGCCCT GTACTTCGTC	660
GCCATGGAGG AGGACGCGG CATGGCCAGC TACCCGAGCA ACCAGGCCGG CGCCCGGTAC	720
GGCACTGGGG TGAGTTGAGC TCCGCTTTGT TTCGAGTCGC AACGAGGCAC TTTCTGGGCG	780
CCGGCTAACT CTCTCGATTC CTCCGACAGT ACTGCGATGC CCAATGCGCT CGTGATCTCA	840
AGTTCGTTGG CGGCAAGGCC AACATTGAGG GCTGGAAGTC GTCCACCAGC GACCCCAACG	900
CTGGCGTCGG CCCGTACGGC AGCTGCTGCG CTGAGATCGA CGTCTGGTGA GTGCGAGACC	960
GTCCACCCAG GTTCGGATGC GGGGTGGAAA TTTCGCGGCT AACGGAGCAC CCCCAGGGA	1020
GTGGAATGCC TATGCCTTCG CTTTACGCC GCACGCGTGC ACGACCAACG AGTACCAGT	1080
CTGCGAGACC ACCAAGTCG GTGGCACCTA CTCGGAGGAC CGCTTCGCCG GCAAGTGCGA	1140
CGCCAACGGC TCGACTACA ACCCCTACCG CATGGGCAAC CCCGACTTCT ACGGCAAGGG	1200
CAAGACGCTC GACACAGCC GCAAGTTTAC GTGCGTGACC CCTTGTGGCG CAACCTTTCT	1260
CTGCCTGCCT GGACACTG AAATGACAC GTCGTTTTCG GCTGCAGCGT CGTCTCCCGC	1320

-continued

TTGAGGAGA ACAAGCTCTC CCAGTACTTC ATCCAGGACG GCCGCAAGAT CGAGATCCCG	1380
CCGCCGACGT GGGAGGGCAT GCCCAACAGC AGCGAGATCA CCCCCGAGCT CTGCTCCACC	1440
ATGTTTCGATG TGTTCACGA CCGCAACCGC TTCGAGGAGG TCGGCGGCTT CGAGCAGCTG	1500
AACAACGCCC TCCGGGTTC CATGGTCCTC GTCATGTCCA TCTGGGACGA CGTAAGTACC	1560
CGCCGACCTC CCTAGCCACA CAAGCCGCAT CCGGCGAGGC ACGCCATCGC TGCTGCTAAC	1620
ACGAGACCGT TCGTAGCACT ACGCCAACAT GCTCTGGCTC GACTCCATCT ACCCGCCCGA	1680
GAAGGAGGGC CAGCCCGGCG CCGCCCGTGG CGACTGCCCC ACGGACTCGG GTGTCCCCGC	1740
CGAGGTCGAG GCTCAGTTCC CCGACGCGTA AGACTTGCCC CCGACCCCAA GCTTCCACTT	1800
CTGGATGCCG AATGCTAACA CGCGAAACAG CCAGGTCGTC TGGTCCAACA TCCGCTTCGG	1860
CCCCATCGGC TCGACCTACG ACTTCTAAGC CGGTCCATGC ACTCGCAGCC TGGGGCCCGT	1920
CACGCCCGCC ACCTCCCCTC GCGGAAACTC TCCGTGCGTC GCGGGCTCCA AAGCATTTTG	1980
GCCTCAAGTT TTTTCGTTT	2000

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 452 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..452
- (D) OTHER INFORMATION: /label= 50K-cellulase-B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

Met Met Met Lys Gln Tyr Leu Gln Tyr Leu Ala Ala Ala Leu Pro Leu	1	5	10	15
Val Gly Leu Ala Ala Gly Gln Arg Ala Gly Asn Glu Thr Pro Glu Asn	20	25	30	
His Pro Pro Leu Thr Trp Gln Arg Cys Thr Ala Pro Gly Asn Cys Gln	35	40	45	
Thr Val Asn Ala Glu Val Val Ile Asp Ala Asn Trp Arg Trp Leu His	50	55	60	
Asp Asp Asn Met Gln Asn Cys Tyr Asp Gly Asn Gln Trp Thr Asn Ala	65	70	75	80
Cys Ser Thr Ala Thr Asp Cys Ala Glu Lys Cys Met Ile Glu Gly Ala	85	90	95	
Gly Asp Tyr Leu Gly Thr Tyr Gly Ala Ser Thr Ser Gly Asp Ala Leu	100	105	110	
Thr Leu Lys Phe Val Thr Lys His Glu Tyr Gly Thr Asn Val Gly Ser	115	120	125	
Arg Phe Tyr Leu Met Asn Gly Pro Asp Lys Tyr Gln Met Phe Asn Leu	130	135	140	
Met Gly Asn Glu Leu Ala Phe Asp Val Asp Leu Ser Thr Val Glu Cys	145	150	155	160
Gly Ile Asn Ser Ala Leu Tyr Phe Val Ala Met Glu Glu Asp Gly Gly	165	170	175	

-continued

Met Ala Ser Tyr Pro Ser Asn Gln Ala Gly Ala Arg Tyr Gly Thr Gly
 180 185 190

Tyr Cys Asp Ala Gln Cys Ala Arg Asp Leu Lys Phe Val Gly Gly Lys
 195 200 205

Ala Asn Ile Glu Gly Trp Lys Ser Ser Thr Ser Asp Pro Asn Ala Gly
 210 215 220

Val Gly Pro Tyr Gly Ser Cys Cys Ala Glu Ile Asp Val Trp Glu Ser
 225 230 235 240

Asn Ala Tyr Ala Phe Ala Phe Thr Pro His Ala Cys Thr Thr Asn Glu
 245 250 255

Tyr His Val Cys Glu Thr Thr Asn Cys Gly Gly Thr Tyr Ser Glu Asp
 260 265 270

Arg Phe Ala Gly Lys Cys Asp Ala Asn Gly Cys Asp Tyr Asn Pro Tyr
 275 280 285

Arg Met Gly Asn Pro Asp Phe Tyr Gly Lys Gly Lys Thr Leu Asp Thr
 290 295 300

Ser Arg Lys Phe Thr Val Val Ser Arg Phe Glu Glu Asn Lys Leu Ser
 305 310 315 320

Gln Tyr Phe Ile Gln Asp Gly Arg Lys Ile Glu Ile Pro Pro Pro Thr
 325 330 335

Trp Glu Gly Met Pro Asn Ser Ser Glu Ile Thr Pro Glu Leu Cys Ser
 340 345 350

Thr Met Phe Asp Val Phe Asn Asp Arg Asn Arg Phe Glu Glu Val Gly
 355 360 365

Gly Phe Glu Gln Leu Asn Asn Ala Leu Arg Val Pro Met Val Leu Val
 370 375 380

Met Ser Ile Trp Asp Asp His Tyr Ala Asn Met Leu Trp Leu Asp Ser
 385 390 395 400

Ile Tyr Pro Pro Glu Lys Glu Gly Gln Pro Gly Ala Ala Arg Gly Asp
 405 410 415

Cys Pro Thr Asp Ser Gly Val Pro Ala Glu Val Glu Ala Gln Phe Pro
 420 425 430

Asp Ala Gln Val Val Trp Ser Asn Ile Arg Phe Gly Pro Ile Gly Ser
 435 440 445

Thr Tyr Asp Phe
 450

(2) INFORMATION FOR SEQ ID NO: 36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 887 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 351..455
- (D) OTHER INFORMATION: /product= "protein-with-CBD"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

CCATGGACGC GAACTGCGAC GTCTTCTGCC CCGAGCTGAA GACCCAGAGC ATCCAGACCG 60

GCAACCACTG CACCCAGGAG ATGAAGGTCT ACGAGAACAT TGACGGCTGG CTCGACAGCC 120

-continued

TGCCCCGGCAA CGTCCCCATC ACCGGTCCGC AGCCCCGGCTC TGGTAAGTCA AAGAGATGAT	180
GCCTACCTAC CTTCACACCT TCCCACCCAG CCGCAAATAC CTTTCTCCCT CCCCCTGCCC	240
CGTATTCTTT CAACGCCCCG AGACTGACAG ACCCGCTCGT CCCAGCGGC AACCCCGGCA	300
ACGGCGGCGG CAGCAACCCG GGCAACGCG GCGGCGGCGG CTGCACCGTC CAGAAGTGGG	360
GCCAGTGGG CGGCATCGGC TACTCGGGCT GCACCACCTG CAAGGCCGGC TCGACCTGCC	420
CGGCCAGAA CGAGTACTAC TCGAGTGCC GTAAAGCGG CCGTGGGCTA GGTGGCCGAG	480
CGGGGGGTT TCTTCATTGG TTGAGCAAT AGAACAGGAT TTCCGGCTCG TTGGCAGCGG	540
CGCGCCGCGG GGATGGTGTT GTACAATTCA AGACCTCAGT ACCGAGGGAC CTGGAAGGA	600
GTGAGTCTGC TTGTACGGAG GCTGGCTGCC CCGTGGCGGC GCTGGCAAGG TAGATAGCCC	660
TTCATTGCTG TAACTAGTAT GCTATATACC TCTGCACATT TGCAGCCCCA TGGTGTGAAC	720
AACAAGTGAC AAGGCTTCCA GTTCCAGCCT CGCGCAATTG TCACGATATC CTGGGTCCAT	780
CTATATGTAT GGGCATGAGC GAGTCGAGAA AATGTACCG GAAAAATCGT AGTGACCTGC	840
GCACTGCGCC GTTCTACCAC CGTAGGATTG AAGTGAATCT CGAATTC	887

(2) INFORMATION FOR SEQ ID NO: 37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Melanocarpus albomyces*
- (B) STRAIN: ALKO4237

(ix) FEATURE:

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..34
- (D) OTHER INFORMATION: /label= prot-with-CBD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

Gln	Lys	Trp	Gly	Gln	Cys	Gly	Gly	Ile	Gly	Tyr	Ser	Gly	Cys	Thr	Thr
1			5					10						15	
Cys	Lys	Ala	Gly	Ser	Thr	Cys	Pro	Ala	Gln	Asn	Glu	Tyr	Tyr	Ser	Gln
			20				25						30		
Cys	Leu														

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

ATAGAATCTCT AYTGGGAYTG YTGAAARCC

29

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

ATAGAATTCT TRTCNGCRTT YTGRAA 26

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GAYGARACNG ARCAIMG 17

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

TANGCNCNC CNGGRTT 17

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

AARCAYGART AYGGNAC 17

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CCRTARAART CNGGRTT 17

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

CCGCGGACTG GCATC

15

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

CCGCGGACTG GCATC

16

What is claimed is:

1. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

2. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having an amino acid sequence which has at least 80% identity to the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

3. An enzyme preparation having cellulase activity, wherein said activity comprises the cellulase activity of a 20K polypeptide having the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31), wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with the nucleic acid sequence set forth in FIG. 19 (SEQ ID NO:30); and
 (ii) separating said host cell from the culture medium and obtaining the supernatant having said 20K cellulase activity;

wherein the level of 20K cellulase activity in the supernatant obtained from said transformed host cell is at least three times the level of 20K cellulase activity obtained from the supernatant of the untransformed host cell naturally expressing said cellulase when said untransformed host cell is cultivated under the same conditions.

4. An enzyme preparation having cellulase activity wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with a nucleic acid sequence encoding a cellulase having at least 80% identity to the amino acid sequence of the 20K cellulase set forth in FIG. 19 (SEQ ID NO:31); and
 (ii) separating said host cell from the culture medium and obtaining the supernatant having said 20K cellulase activity or cellulase activity of a cellulase that has at least 80% identity to the sequence of said 20K cellulase set forth in FIG. 19 (SEQ ID NO:31),

wherein the level of said 20K cellulase activity or the level of said activity of said cellulase that has at least 80% identity to said sequence of said 20K cellulase set forth in FIG. 19 (SEQ ID NO: 31) in the supernatant obtained from said transformed host cell is at least three times the level of said 20K cellulase activity or activity of said cellulase that has at least 80% identity to said sequence of said 20K cellulase set forth in FIG. 19 (SEQ ID NO:31), respectively, obtained from the supernatant of the strain naturally expressing said cellulase when said strain is cultivated under the same conditions.

5. An enzyme preparation comprising at least one cellulase from *Melanocarpus albomyces* CBS 685.95.

6. The enzyme preparation of any one of claims 3-5, which is in the form of a liquid.

7. The enzyme preparation of any one of claims 3-5, which is in the form of a dry powder.

8. A detergent composition comprising the enzyme preparation of any one of claims 3-5 in combination with a surface active agent or surfactant.

9. A method of treating cellulosic fiber containing textile material, said method comprising contacting said textile material with the detergent composition of claim 8.

10. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

11. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having an amino acid sequence which has at least 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

12. An enzyme preparation having cellulase activity, wherein said activity comprises the cellulase activity of a polypeptide having amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31), wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with a nucleic acid sequence encoding amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31); and

(ii) separating said host cell from the culture medium and obtaining the supernatant having said cellulase activity; wherein the level of said cellulase activity in the supernatant obtained from said transformed host cell is at least three times the level of said cellulase activity obtained from the supernatant of the strain naturally expressing said cellulase when said strain is cultivated under the same conditions.

13. An enzyme preparation having cellulase activity wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with a nucleic acid sequence encoding a cellulase having at least 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31); and
 (ii) separating said host cell from the culture medium and obtaining the supernatant having the cellulase activity of a cellulase that has at least 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31);

95

wherein the level of said activity of said cellulase that has at least 80% identity to amino acids 22–235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31) in the supernatant obtained from said transformed host cell is at least three times the level of said activity of said cellulase 5 that has 80% identity to amino acids 22–235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31) obtained from the supernatant of the strain naturally expressing said cellulase when said strain is cultivated under the same conditions.

96

14. The enzyme preparation of either claim 12, or claim 13, which is in the form of a liquid.

15. The enzyme preparation of either claim 12 or claim 13, which is in the form of a dry powder.

16. A detergent composition comprising the enzyme preparation of either claim 12 or claim 13 in combination with a surface active agent or surfactant.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,723,549 B2
DATED : April 20, 2004
INVENTOR(S) : Miettinen-Oinonen et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title page.

Item [75], Inventors, please delete the list and insert the following corrected list:

-- Arja Miettinen-Oinonen, Masala (FI);
John Londesborough, Helsinki (FI);
Jari Vehmaanperä, Klaukkala (FI);
Heli Haakana, Espoo (FI);
Arja Mäntylä, Helsinki (FI);
Raija Lantto, Klaukkala (FI);
Marja Paloheimo, Vantaa (FI);
Pirkko Suominen, Helsinki (FI) --.

Column 5.

Line 20, please delete "FIG. 19" and insert therefore -- FIGS. 19 (A and B) --.

Signed and Sealed this

Twenty-seventh Day of July, 2004

A handwritten signature in black ink, appearing to read "Jon W. Dudas", written over a horizontal line.

JON W. DUDAS
Acting Director of the United States Patent and Trademark Office